



## **SURAT KETERANGAN**

**Nomor: 62/XI/Adm-P-P3M/SHT/2022**

Pusat Penelitian, Pengembangan dan Pengabdian Kepada Masyarakat (Pusat P3M) Stikes Hang Tuah Surabaya menerangkan bahwa telah selesai melaksanakan pemeriksaan plagiarisme dengan membandingkan artikel-artikel lain menggunakan perangkat lunak melalui <https://www.turnitin.com/> pada tanggal 24 Agustus 2022.

Penulis : Ninik Ambar Sari, Ceria Nurhayati  
Judul : *The Relationship Between Self-Management and Quality Of Life Among Patients with Type 2 Diabetes Mellitus*  
No. Pemeriksaan : 1886251888.2022.08.24

Dengan hasil sebagai berikut:

**Tingkat kesamaan di seluruh artikel (*Similarity Index*) sebesar 18%**

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya.

Surabaya, 24 November 2022  
Kepala Pusat P3M  
STIKES Hang Tuah Surabaya



**Christina Yuliastuti, S.Kep., Ns., M.Kep.**  
NIP.03.017

# THE RELATIONSHIP BETWEEN SELF-MANAGEMENT AND QUALITY OF LIFE AMONG PATIENTS WITH TYPE 2 DIABETES MELLITUS

*by Ninik Ambar Sari*

---

**Submission date:** 24-Aug-2022 11:07AM (UTC+0700)

**Submission ID:** 1886251888

**File name:** The\_Relationship\_Between\_Self\_Management.pdf (152.23K)

**Word count:** 3445

**Character count:** 18592

Original Research Article

7

## THE RELATIONSHIP BETWEEN SELF-MANAGEMENT AND QUALITY OF LIFE AMONG PATIENTS WITH TYPE 2 DIABETES MELLITUS

Ninik Ambar Sari<sup>1\*</sup>, Ceria Nurhayati<sup>1</sup>

<sup>1</sup> STIKES Hang Tuah Surabaya, East Java Province, Indonesia

\*Correspondence:

Ninik Ambar Sari

STIKES Hang Tuah Surabaya, East Java Province, Indonesia  
Julia Gadung No. 1, Surabaya Regency,  
East Java Province, Indonesia

Email:

ninik@stikeshangtuah.ac.id

Article Info:

Received: August 6, 2021

Revised: December 27, 2021

Accepted: December 29, 2021

DOI:

<https://doi.org/10.36720/njk.v10i2.300>

Abstract

13

**Background:** Diabetes Mellitus (DM) is a metabolic disease characterized by high blood glucose levels as a result of insufficient insulin secretion, impaired insulin activity or both. Patients with diabetes mellitus are at risk for complications that can affect their quality of life. These complications can be minimized through self-management.

**Objectives:** The aim of this study was to determine the relationship between self-management and quality of life in type 2 DM patients.

**Design:** This study was a quantitative research. The design in this study was a cross-sectional analytic with a sample of 75 people who were carried out at the endocrine polyclinic RS Pusat TNI Angkatan Laut (RSPAL) DR. Rangga Surabaya. Research instrument used the Diabetes Self-Management Questionnaire (DSMQ) developed by Schmitt et al (2013) to determine the self-management of type 2 DM patients and to measure the quality of life using the Diabetes Quality of Life Brief (DQoL-Brief) instrument developed by Burrough et al (2008). The sample size with simple random sampling method, and data analysis using spearman rank test.

**Results:** The results showed that there was a significant relationship between self-management and quality of life in type 2 DM patients ( $p$ -value = .000,  $r_s = .394$ ).

**Conclusion:** There is a significant positive relationship between the level of self-management and quality of life in type 2 DM patients. From the results of the Spearman test correlation coefficient, there is a positive  $r_s = .141$ . This means that the higher the patient's self-management, it will also improve the quality of their life. The implication of this study are nurses can increase patient knowledge by providing education that focuses on improving self-management and facilitating the provision of family support as well as supervision and monitoring related to self-management by type 2 DM patients.

**Keywords:** Self-management, quality of life, type 2 diabetes mellitus.

1

© 2021 The Author. Nurse and Health: Jurnal Kependidikan Published by Institute for Research and Community Service

E-ISSN

2623-2448

Health Polytechnic of Kerta Cendekia, Surabaya

P-ISSN

2088-9909

This is an Open Access Article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 IGO (CC BY-NC 4.0), which allows others to remix, tweak, and build upon the work non-commercially as long as the original work is properly cited. The new creation are not necessarily licensed under the identical terms.

### INTRODUCTION

Diabetes mellitus is a group of metabolic disorders characterized by chronic

hyperglycemia, which is caused by impaired insulin secretion, insulin action or both (ADA, 2012). The International Diabetes Federation

4

(IDF) states that diabetes mellitus is a major health problem. The main problem faced by patients with diabetes mellitus is chronic hyperglycemia which can lead to various complications (Papatheodoro, et al., 2016). The results of a study conducted on 66,726 patients with type 2 diabetes mellitus proved that 53% had microvascular complications and 27.2% had macrovascular complications. (Litwak et al., 2013).

Diabetes mellitus can be prevented or its occurrence can be delayed and controlled, with optimum treatment management (Brady, et al., 2013). Where the higher the quality of life of patients with diabetes mellitus, the prevention of diabetes mellitus is also good, but in fact a decrease in the quality of life in patients with diabetes mellitus is often followed by the inability of these patients to carry out self-management independently (Choi, et al., 2011). The inability of patients with diabetes mellitus to perform self-management can affect the quality of life in terms of physical health, psychological well-being, social relationships, and relationships with the environment (Joukman, et al., 2016). In this case, quality of life is an important concern for health professionals because it can be a reference for the success of an action or therapy. Diabetes mellitus will accompany the patient's lifetime so that it greatly affects a person's quality of life. Low quality of life can exacerbate complications and can end in disability or death (Brady, et al., 2013).

One of the efforts that can be done to improve diabetes outcomes in diabetes management is diabetes self-management (DSM). The results of a systematic review conducted on 3,421 research articles also prove that diabetes self-management is effective in supporting a person's success in managing diabetes (Heinrich, Schaper, & Vries, 2010). The level of knowledge about DM affects the patient's ability to perform self-management. In addition, knowledge of DM can be used as a basis for decision making about diet, exercise, blood glucose monitoring, drug use, weight control, and foot care. (Palmer, et al., 2014). The ability to take care of yourself is not only

determined by attitude. However, attitudes and abilities in good self-management and monitoring the daily lifestyle behavior of DM patients and changing old habits are one of the efforts to control diabetes. Thus, the factor of understanding or knowledge about diabetes with one's self-management behavior is very important to be considered by health professionals. Attitude is an important role of emotional response and affects the patient's efforts to manage diabetes in daily life (Perkeni, 2015). Self-management of diabetes mellitus can develop patient skills in problem solving, increase self-confidence and support the application of knowledge in real life (Choi, et al., 2011).

Based on the results of interviews with 5 people with diabetes mellitus at RSPAL DR. Ramlan Surabaya, showed that 3 of them experienced low quality of life due to fatigue during activities and disturbed sleep. One of them said that he only took medication if there were complaints and an inability to follow a diet. Based on this description, the researcher is interested in describing the relationship between self-care and quality of life in type 2 diabetes mellitus patients at the endocrine polyclinic RS Pusat TNI Angkatan Laut (RSPAL) DR. Ramlan Surabaya

## METHODS

### Study design

This research is a quantitative study with a descriptive correlation method through a cross sectional approach.

### Setting

This research was conducted at the endocrine polyclinic RS Pusat TNI Angkatan Laut (RSPAL) DR. Ramlan Surabaya.

### Research Subject

This research was conducted from 20 February to 20 March 2021. The population in this study were all patients suffering from type 2 diabetes mellitus. The sample size in this study was 75 patients. The sampling technique in this study used simple random sampling method. The sample is then selected based on

the characteristics and criteria of the sample based on:

1. Inclusion Criteria
  - a. Patients who come to the endocrine polyclinic RSPAL Surabaya
  - b. Patients who have been treated at the endocrine polyclinic RSPAL
  - c. Diagnosed with type 2 diabetes mellitus.
  - d. Respondent in the study fill out the questionnaire given.
  - e. Cooperative.
2. Exclusion Criteria
  - a. Diagnosed with type 1 diabetes mellitus.
  - b. Patients who cannot remember activities during the previous 1 week.

#### *Instrument*

The instrument used is the Diabetes Self-Management Questionnaire (DSMQ) developed by Schmitt et al (2013) to determine the self-management of type 2 DM patients. The DSMQ questionnaire instrument that has been analyzed using the validity and reliability test of the SPSS 24 computerized system for windows with degrees of freedom  $30-2 = 28$  ( $r$  table = .374) gets the results from 16 statements that have 2 invalid statements, namely number 4 ( $r = -.171$ ) and number 9 ( $r = -.044$ ), but because the substance of the question is considered important, the question is not deleted but the statement is corrected. The DSMQ questionnaire is reliable, that is, by using the alpha test, the Cronbach alpha value is .741. The instrument for quality of life using

used to the Diabetes Quality of Life Brief (DQoL Brief) instrument developed by Burrough et al. (2008). The diabetes quality of life questionnaire is reliable, that is, by using the alpha test, the Cronbach alpha value is .963.

#### *Data Analysis*

After the data is collected the researcher sorts out the research data and does the coding. Data analysis using SPSS version 25. The analysis used the Spearman rank test with a significance level  $\alpha < .05$ .

#### *Ethical Consideration*

This research has been reviewed and decided to have passed the ethical review of the RS Pusat TNI Angkatan Laut (RSPAL) DR. Ramlan Surabaya Ethics Committee. Ramlan Surabaya with number SKET/3348/II/2021 in an effort to protect the human rights and welfare of nursing research subjects. At the time of conducting the research, the researcher guaranteed all the confidentiality of the respondents and did not violate the rights of the respondents and did not cause harm to the respondents who participated in this research.

## RESULTS

#### *Characteristics of Respondents*

The results of data analysis about the characteristics of the respondents (table 1) showed that the majority were male as many as 39 respondents (52%). The characteristics of the respondents' age showed that most of them were 56-75 years old as many as 40 respondents (53%), and the proportion of good HbA1c values is 46 respondents (61.3%).

Table 1. Distribution of Frequency of Respondents in the Endocrine Polyclinic RS Pusat TNI Angkatan Laut (RSPAL) DR. Ramlan Surabaya from February 20, 2021 to March 20, 2021.

Characteristics of Respondents	Frequency (f)	Percentage (%)
<i>Gender</i>		
Male	36	48.0
Female	39	52.0
<i>Age (Years)</i>		
40-55	32	42.7
56-75	40	53.3

Characteristics of Respondents	Frequency (f)	Percentage (%)
> 75	3	4.0
Value of HbA1c		
Bad	29	38.7
Good	46	61.3

Source: Primary Data of Questionnaire, 2021.

#### Analysis of <sup>10</sup> Correlation between Self-Management and Quality of Life among Patients Type 2 Diabetes Mellitus

The result data on <sup>11</sup> table 2 showed that self-management had relationship with quality of

life among patients type 2 diabetes mellitus positively ( $p$ -value = .000,  $r$  = .349). This result mean relationship between self-management and quality of life was good enough.

**Table 2.** Relationship between Self-Management and Quality of Life among Patients Type 2 Diabetes Mellitus in the Endocrine Polyclinic RS Pusat TNI Angkatan Laut (RSPAL) DR. Ramaela Surabaya from February 20, 2021 to March 20, 2021.

Self-Management	Quality of Life						Total
	Not Good		Sufficient		Good		
f	%	f	%	f	%	f	%
Not Good	11	14.7	19	25.3	0	0.0	30 40.0
Sufficient	0	0.0	25	33.3	10	13.3	35 16.7
Good	0	0.0	1	1.4	9	12.0	10 13.3
Total	11	14.7	45	60.0	19	25.3	75 100.0

<sup>11</sup>  $p$ -value = .000;  $r$  = .349

Source: Primary Data of Questionnaire, 2021.

#### DISCUSSION

The results showed that most of the respondents were aged 56-75 years with a total of 40 respondents (53%). Based on the pathogenesis of DM, the aging process is at risk for the occurrence of DM or called prediabetes, namely the condition of the patient's blood sugar having impaired tolerance (140-199 mg/dL). If the blood sugar level reaches 200 mg/dL then it is classified as diabetes mellitus. Wulansari & Isfandiari (2013) stated that increasing age resulted in the ability of insulin as the key to enter glucose into cells not functioning properly, resulting in insulin resistance.

Naturally, as a person ages, changes occur both physiologically, psychologically and intellectually. The addition of age, especially in the elderly, will have an impact on anatomical, physiological and biochemical change <sup>12</sup>. This will result in susceptibility to a disease and can

lead to failure to maintain homeostasis against stress. Failure to maintain this homeostasis will result in a decrease in the body's resistance to life and result in an increase in the ease of emergence of several disorders in the individual (Hayes, et al., 2013). Whereas in type 2 diabetes patients, the changes that occur will have an impact on increasing impaired glucose tolerance and insulin resistance. This will have an impact on various problems both physiologically, psychologically and socially. <sup>13</sup> This will cause various limitations that will affect the quality of life. In addition to the age factor and decreased body function, it also has an impact on the decreased ability in self-care and the implementation of diabetes management so that health problems will easily arise. This of course will affect the quality of life (Choi, et al., 2011).

The next respondent characteristic is that most of the sufferers are women as many as 39

respondents (52%). Women are more at risk of developing DM because physically women have a greater chance than men to develop central obesity and increase body mass index (Yusra, 2011). <sup>12</sup>

Self-management has a relationship with quality of life, this is in accordance with table 2 statistical test results value <0.05 with a positive correlation direction, which means that if self-management is carried out regularly, the quality of life of DM sufferers will increase. This is in line with the results of a systematic review study to see the relationship between diabetes self-management and quality of life in 7,878 literatures conducted from 1985 to 2013 proving a significant relationship that self-management can improve the quality of life of patients with chronic diseases, namely diabetes (Jonkman et al., 2016). In line with the meta-analysis research conducted on 23 research articles in determining the effectiveness of the Chronic Disease Self-Management Program (CDSMP) it proved that it was effective in improving behavior change, physical and psychological health. Changes in health behavior including physical activity, cognitive management and communication with doctors increased significantly so that the quality of life improved (Brady et al., 2013). In line with the results of a meta-analysis conducted on 1,892 subjects to see a comparison between the intervention group given diabetes self-management training and the control group, the results showed that there was an increase in the value of quality of life in the intervention group while there was no increase in the control group (Choi, Y J, et al., 2011).

Based on the researcher's assumption that self-management is one of the strongest predictors of quality of life. However, the contribution of various other factors also needs to be considered. Family support, as well as high education and avoiding various complications will certainly further improve the health of type 2 diabetes patients, so that components of quality of life such as physical, psychological and social functions will

certainly be well maintained (Brandy, T.J, et al., 2013).

Based on the assumption of the researcher, naturally along with the age of a person changes occur both physiologically, psychologically and intellectually. The addition of age, especially in the elderly, will have an impact on anatomical, physiological and biochemical changes. <sup>13</sup> will result in susceptibility to a disease and can lead to failure to maintain homeostasis against stress. Failure to maintain this homeostasis will result in a decrease in the body's resistance to life and result in an increase in the case of emergence of several disorders in the individual (Zychowska, et al., 2013).

## CONCLUSION

There is a significant positive relationship between the level of self-management and quality of life in type 2 DM patients. From the results of the Spearman test correlation coefficient, there is a positive sign. This means that the higher the patient's self-management, it will also improve the quality of their life.

## SUGGESTION

The implication of this study are nurses can increase patient knowledge by providing education that focuses on improving self-management and facilitating the provision of family support as well as supervision and monitoring related to self-management by type 2 DM patients.

## ACKNOWLEDGMENT

The authors express their gratitude of STIKES Hang Tuah Surabaya and RS Pusat TNI Ang 10 in Laat (RSPAL) DR. Ramelan Surabaya who has given their permission for the authors conduct research, guided the authors, and to all the participants who were involved.

## DECLARATION OF CONFLICTING INTEREST

There is no conflict of interest in the writing.

## FUNDING

This research was carried out using funding from the author himself.

## AUTHOR CONTRIBUTION

**Ninik Ambar Sari:** Contributes to the completion of the article.

**Ceria Nurhayati:** Contributes to the completion of the article.

## ORCID

**Ninik Ambar Sari:**

<https://orcid.org/0000-0003-2456-2107>

**Ceria Nurhayati:**

None.

## REFERENCES

- ADA. (2020). Standards of medical care in diabetes-2012. *Diabetes Care*, 43 (SUPPL.1). <https://doi.org/10.2337/dc12-SINT>
- Brady, T. J., Murphy, L., Colmain, B. J. O., Beauchesne, D., Daniels, B., Greenberg, M., Chervin, D. (2013). A Meta analysis of health status, health behaviors, and health care utilization outcomes of the chronic disease self management program. *Preventing Chronic Disease*, 10, 1-14. <https://doi.org/10.5888/pcd10.120112>
- Burroughs, T. E., Desikan, R., Waterman, B. M., Gilin, D., & Megill, J. (2004). Development and Validation of the Diabetes Quality of Life Brief Clinical Inventory. *Diabetes Spectrum*, 17 (1), 41-49.
- Choi, Y.J., Lee, M.S., An, S.Y., Kim, T.H., Han, S.J., Chung, Y.S., Lee, K.W & Kim, D.J., 2011. The Relationship between Diabetes Mellitus and Health-Related Quality of Life in Korean Adults: The Fourth Korea National Health and Nutrition Examination Survey (2007-2009). *Diabetes Metabolism Journal*, 35(6), pp. 587-594.
- Hayes, A. J., Leal, J., Gray, A. M., Holman, R. R., & Clarke, P. M. (2013). UKPDS outcomes model 2: a new version of a model to simulate lifetime health outcomes of patients with type 2 diabetes mellitus using data from the 30 year United Kingdom Prospective Diabetes Study: UKPDS 82 Lipids in Diabetes Study. *Diabetologia*, 56, 1925-1933. <http://doi.org/10.1007/s00125-013-2940-y>
- Heinrich, E., Schaper, N. C., & Vries, N. K. De. (2010). Self management interventions for type 2 diabetes: a systematic review. *EDN Autumn*, 7 (2).
- Jonkman, N. H., Schuurmans, M. J., Groenwald, R. H. H., Hoes, A. W., & Trappenburg, J. C. A. (2016). Identifying components of self management interventions that improve health related quality of life in chronically ill patient: systematic review and meta regression analysis. *Journal Patient Education and Counseling*, 99 (7), 1087-1098. <http://doi.org/10.1016/j.jpec.2016.01.022>
- Litwak, L., Goh, S.-Y., Hassein, Z., Malek, R., Prusty, V., & Khamseh, M. E. (2013). Prevalence of diabetes complications in people with type 2 diabetes mellitus and its association with baseline characteristics in the multinational A1chieve study. *Diabetology & Metabolic Syndrome*, 5 (1), 57. <https://doi.org/10.1186/1758-5996-5-57>
- Palmer, J. S., Brandle, M., Trevisan, R., Federici, M. O., Liabai, S., & Valentine, W. (2014). Assessment of the association between glycemic variability and diabetes-related complications in type 1 and type 2 diabetes. *Journal Diabetes Research and Clinical Practice*, 105, 273-284. <http://doi.org/10.1>
- Papatheodoreou, K., Papanas, N., Banach, M., papazoglou, D., & Edmonds, M. (2016). *Complications of Diabetes 2016, 2016*
- Perkeni. (2015). *Konsensus Pengelolaan dan Pencegahan Diabetes Mellitus Tipe 2 di Indonesia*. PB Perkeni
- Schmitt, A., Gahr, A., Hermanns, N., Kulzer, B., Huber, J., & Haak, T. (2013). The

- Diabetes Self-Management Questionnaire (DSMQ): development and evaluation of an instrument to assess diabetes self-care activities associated with glycaemic control. *Health and Quality of Life Outcomes*, 11, 1. <http://doi.org/10.1186/1477-7525-11-138>
- Yusra, A. (2011). *Hubungan antara dukungan keluarga dengan kualitas hidup pasien diabetes mellitus tipe 2 di poliklinik penyakit dalam rumah sakit umum pusat fatmawati jakarta*. <http://eprints.ui.ac.id>
- Wulandari, Y.M & Isfandiari, A.M. (2013). Kaitan Sindroma Metabolik dan Gaya Hidup dengan Gejala Komplikasi Mitravaskuler. *Jurnal Berkala Epidemiologi*, 1(2)
- Zychowska, M., Rojewska, E., Przewlocka, B., & Mika, J. (2013). Mechanisms and pharmacology of diabetic neuropathy—experimental and clinical studies. *Pharmacological Reports*; PR, 65(6), 1601-10. [https://doi.org/10.1016/S1734-1140\(13\)71521-4](https://doi.org/10.1016/S1734-1140(13)71521-4)

**Cite this article as:** Sari, N. A., Nurhayati, C. (2021). The relationship between self-management and quality of life among patients with type 2 diabetes mellitus. *Nurse and Health: Jurnal Keperawatan*, 10 (2), 343-349. <https://doi.org/10.36720/nhjk.v10i2.300>

# THE RELATIONSHIP BETWEEN SELF-MANAGEMENT AND QUALITY OF LIFE AMONG PATIENTS WITH TYPE 2 DIABETES MELLITUS

---

ORIGINALITY REPORT

---



PRIMARY SOURCES

---

- 1 Dini Prastyo Wijayanti, Riesmiyatiningdyah  
Riesmiyatiningdyah, Kusuma Wijaya Ridi  
Putra, Agus Sulistyowati et al.  
"SOCIALIZATION OF COVID-19 MANAGEMENT  
IN HEALTH POLYTECHNIC OF KERTA  
CENDEKIA", Community Service Journal of  
Indonesia, 2022  
Publication
- 2 Livana P. H., Indah Permata Sari, Indah  
Hermanto. "Description of the Level of  
Ansietas of Patients of Diabetes Mellitusin  
Kendal District", International Journal of  
Diabetes and Endocrinology, 2019  
Publication
- 3 WWW.MDPI.COM  
Internet Source
- 4 Rahayu Budi Utami, Amrina Rosada.  
"PARENTING PATTERNS OF PARENTS AND  
MENTAL-EMOTIONAL DEVELOPMENT IN

TAMAN KANAK-KANAK DHARMA WANITA  
GODEAN, NGANJUK DISTRICT", Nurse and  
Health: Jurnal Keperawatan, 2019

Publication

---

- 5 Samira Mokhlesi, Masoumeh Simbar,  
Fahimeh Ramezani Tehrani, Nourossadat  
Kariman, Hamid Alavi Majd. "Quality Of Life  
Questionnaire for women with Gestational  
Diabetes Mellitus (GDMQ-36): Development  
and Psychometric Properties", Research  
Square Platform LLC, 2019  
Publication
- 6 scholarworks.uttyler.edu 1 %  
Internet Source
- 7 Submitted to University of Lincoln 1 %  
Student Paper
- 8 Yoga Kertapati. "Tugas Kesehatan Keluarga  
dan Tingkat Kemandirian Keluarga di Wilayah  
Pesisir Kota Surabaya", Jurnal Ilmiah  
Keperawatan Stikes Hang Tuah Surbaya, 2019  
Publication
- 9 researchbank.acu.edu.au 1 %  
Internet Source
- 10 Submitted to Forum Perpustakaan Perguruan  
Tinggi Indonesia Jawa Timur 1 %  
Student Paper
-

- 11 Titiek Idayanti, Kurnia Indriyanti, Widya Anggraeni. "THE EFFECT OF EARLY BREASTFEEDING INITIATION (IMD) ON NEW BORN BABY TEMPERATURE IN BPS HEPPY RINA, S.ST, SEDURI VILLAGE-MOJOSARI AND BPS FIFIT, S.ST, PANJER VILLAGE-MOJOSARI", Nurse and Health: Jurnal Keperawatan, 2019  
Publication
- 
- 12 [revistaschilenas.uchile.cl](http://revistaschilenas.uchile.cl) 1 %  
Internet Source
- 
- 13 Submitted to Manipal University 1 %  
Student Paper
- 
- 14 [etd.uwc.ac.za](http://etd.uwc.ac.za) 1 %  
Internet Source
- 
- 15 [search.webdepozit.sk](http://search.webdepozit.sk) 1 %  
Internet Source
- 
- 16 Mohammad Farris Iman Leong Bin Abdullah, Sy-Cherng Luke Woon, Hatta Sidi, Arun Ravindran et al. "Depression, anxiety, and associated factors in patients with diabetes: evidence from the Anxiety, Depression, and Personality Traits in Diabetes Mellitus (ADAPT-DM) study", Research Square Platform LLC, 2020  
Publication
- 
- 17 [journal.ukmc.ac.id](http://journal.ukmc.ac.id) 1 %  
Internet Source

---

18

repo.poltekkesdepkes-sby.ac.id

Internet Source

1 %

---

Exclude quotes Off

Exclude bibliography On

Exclude matches < 1%