



Scopus®

MJN

**THE MALAYSIAN
JOURNAL OF NURSING**

October 2022 | Volume - 14 | Issue - 2

Print - ISSN : 2231-7007 | Online - ISSN - 2462-246X

KDN PP 15780/01/2013 (031712)

ARTICLES

MESSAGE FROM THE EDITOR-IN-CHIEF

Datuk Prof. Dr. Hjh. Bibi Florina Abdullah

2-2

 HTML

 PDF

ORIGINAL ARTICLE

Pre-Operative Patient's Anxiety Level with RT-PCR Swab Covid-19 in Operating Room RSPAL Dr. Ramelan Surabaya

Harun Pabendon, Sri Anik Rustini Rustini, Merina Widyastuti, Ninik Ambar Sari

3-6

 HTML

 PDF

 DOI: [HTTPS://DOI.ORG/10.31674/MJN.2022.V14I02.001](https://doi.org/10.31674/mjn.2022.v14i02.001)

Association Between Nutritional Habits and Glycosylated Hemoglobin Level in Type 2 Diabetes Mellitus

Salwa Hagag Abdelaziz, Ola Ibrahim Abdo, Seham Kamal Abde Llatif

7-13

 HTML

 PDF

 DOI: [HTTPS://DOI.ORG/10.31674/MJN.2022.V14I01.002](https://doi.org/10.31674/mjn.2022.v14i01.002)

Relationship between Diet Patterns and the Incidence of Anemia among Adolescent Girls at SMA Giki 1 Surabaya

Novan Hariyanto, Iis Fatimawati, Puji Hastuti, Astrida Budiarti, Sandeep Poddar

14-19

Neurologic Examination of the Diabetic Foot to Prevent Diabetic Foot Ulcer at Puskesmas Kebonsari Surabaya, Indonesia

Nur Muji Astuti, Imroatul Farida, Nabelatul Laili Fauzia, Ceria Nurhayati, Nuh Huda, Didik Winarno, Ruma Poddar

90-95



HTML



PDF



DOI: [HTTPS://DOI.ORG/10.31674/MJN.2022.V14I02.015](https://doi.org/10.31674/mjn.2022.v14i02.015)

Family Support Related to Quality of Life in Diabetes Mellitus Patients with Ulcer

Christina Yuliasuti, Nur Muji Astuti, Didik Dwi Winarno, Mei Ayu Sari, Norin John

96-100



HTML



PDF



DOI: [HTTPS://DOI.ORG/10.31674/MJN.2022.V14I02.016](https://doi.org/10.31674/mjn.2022.v14i02.016)

The Role of Psychological Distress in Relational Health Instability among Romantic Couples: A Qualitative Inquiry

Michelle A. Calda-Tolibas

101-108



HTML



PDF



DOI: [HTTPS://DOI.ORG/10.31674/MJN.2022.V14I02.017](https://doi.org/10.31674/mjn.2022.v14i02.017)

Psychological Impact of Coping Strategies and Nurse Performance During the Covid-19 Pandemic at Rspal Dr. Ramelan Surabaya

Setiadi, Dwi Priyantini, Dedi Irawandi, Sandeep Poddar

109-116

Neurologic Examination of the Diabetic Foot to Prevent Diabetic Foot Ulcer at Puskesmas Kebonsari Surabaya, Indonesia

Nur Muji Astuti¹, Imroatul Farida^{1*}, Nabelatul Laili Fauzia¹, Ceria Nurhayati¹, Nuh Huda¹, Didik Winarno², Ruma Poddar³

¹Sekolah Tinggi Ilmu Kesehatan Hang Tuah Surabaya, Jawa Timur 60244, Indonesia

²Rumah Luka Surabaya, East Java 60244, Indonesia

³Lincoln University College, 47301 Petaling Jaya, Malaysia

*Corresponding Author's Email: imroatulfarida@stikeshangtuah-sby.ac.id

ABSTRACT

Background: Neuropathy is a complication of DM that changes the structure and function of peripheral nerves, causing diabetic neuropathy due to peripheral nerve degeneration. Neuropathy examination is used to prevent Diabetic Foot Ulcers (DFU), but it is still rarely done or has not become a routine examination in health services. Alternative examination of neuropathy with the Ipswich Touch Test (IpTT). The aim of this study was to identify neurosensory-peripheral disorders in the diabetic foot. **Methods:** The research design used a quantitative description with a cross-sectional approach involving 40 respondents. Intervention with diabetic foot examination using the Ipswich Touch Test (IpTT) instrument which the neurosensory peripheral measurement decreased if the IpTT score was below 2-6. Sampling technique with accidental sampling approach. **Results:** The results of the neurosensory peripheral examination in patients with type 2 diabetes mellitus through the IpTT examination had an average value of 3.85 out of a total score of 6, which means that the respondent experienced a decrease in sensory function. **Conclusion:** The majority of patients with diabetes mellitus type 2 have decreased neurosensory peripheral function, so it will be high risk of diabetic foot ulcers. Nurses can perform an Ipswich Touch Test (IpTT) on diabetic feet to assess peripheral neurosensory function to prevent diabetic foot ulcers.

Keywords: Neuropathy; Neurosensoriperipheral; Ipswich Touch Test (IpTT); Diabetes Mellitus

INTRODUCTION

Neuropathy is a change in the structure and function of the motoric, sensory, and autonomic peripheral nerves, which causes diabetic neuropathy due to peripheral nerve degeneration (Sembiring, Simbolon & Lase, 2018). Neuropathy causes pain, tingling, numbness or numbness, numbness, muscle stiffness, cramps, hypersensitivity to impaired bladder control, and muscle weakness. The problem of neuropathy in people with diabetes mellitus is also exacerbated by a decrease in the immune system so that they are susceptible to infection. Patients with diabetes mellitus (DM) who experience the slightest wound will be very easy to experience tissue necrosis which ends in amputation if not handled properly (Sembiring, Simbolon & Lase, 2018). People with diabetes may develop several different foot problems as a result of nerve damage (Abo Deif & Abdelaziz, 2019). Neurological examination to prevent Diabetic Foot Ulcers (DFU) in patients with type 2 diabetes mellitus is

still rarely done or has not become a routine examination at the Kebonsari Health Center Surabaya. The reality in the field is that there are still many people with diabetes mellitus who experience neurological disorders of the feet, both micropathies, and macropathies, who have not had a neurological examination to reduce the risk of DFU. Patient's autonomy rises in importance, it becomes increasingly important for patients to assume responsibility for their health care (Rosli *et al.*, 2022).

According to The Foundation for Peripheral Neuropathy, DM sufferers who experience peripheral neuropathy are as much as 25% of DM patients in the world (Hicks & Selvin, 2019). Peripheral neuropathy is the most common complication in DM patients and 50% of type 2 DM patients (Tabatabaei-Malazy *et al.*, 2011). The PERSI and Information Center (Indonesian Hospital Association) stated that the prevalence of neuropathy in 2011 in DM patients in Indonesia was more than 50%. This statement is reinforced by the Basic Health Research (Risksedas, 2018) which shows that the most

Received June 29, 2022; Received in revised form September 22, 2022; Accepted September 29, 2022

diabetes complications are neuropathy. BPJS patients data in 2016 stated that DM patients who experienced chronic complications of neuropathy were recorded at the Internal Medicine Poly Hospital of Dr. M. Soewandhi was still quite high in 2016 as many as 691 people (71.90%) of which 497 people were caused by type 2 DM (Putri, 2015). The results of a preliminary study at the Kebonsari Public Health Center in Surabaya found that out of 10 patients with diabetes mellitus type 2 were examined, there were 8 people (80%) with neurological disorders in diabetic feet and 2 people (20%) with normal diabetic feet.

DM with neuropathy will increase the risk of DFU seven times higher than DM patients without neuropathy (Soheilykhah *et al.*, 2018). Patients with DM if blood glucose levels are not controlled will occur chronic complications in the form of nerve disorders called neuropathy. Neuropathy causes changes in nerve tissue due to the accumulation of sorbitol and fructose resulting in loss of axons, decreased speed of induction, paresthesias, decreased muscle reflexes, muscle atrophy, excessive sweating, dry skin, and loss of taste, accelerate the occurrence of diabetic foot ulcers (Alport & Sander, 2012).

Efforts to detect neuropathic disorders in diabetic feet to prevent diabetic foot ulcers need a neurological examination. A neurological examination can be done through sensitivity examination using the Ipswich Touch Test (IpTT). IpTT is a method for early detection of Diabetic Foot ulcers (DFU) that is easy to do, fast, effective, does not cost money, is sensitive, specific, and does not require other instruments (Sharma *et al.*, 2014). How to use IpTT by touching the tips of the first, third, and fifth toes using the examiner's finger for 1-2 seconds. The touch should be light and gentle. If the patient does not feel the touch 2 points then it is declared neuropathy (Diabetes, 2012). The purpose of this study was to identify peripheral neurosensory disorders in the diabetic foot.

METHODOLOGY

The research design used descriptive quantitative with a cross-sectional approach. The study was conducted in May 2020. The population in this study was 45 people with diabetes mellitus type 2 at Puskesmas Kebonsari Surabaya. The sample of this study amounted to 40 people using accidental sampling. The population data questionnaire was used to examine the population data of respondents which included gender, age,

education, work history, cigarette consumption, diabetic foot exercise, taking DM medication, and blood sugar control. Intervention for neurosensory peripheral examination of the diabetic foot using the Ipswich Touch Test (IpTT) instrument (Basir *et al.*, 2020; McLaren & Lu, 2022; Sulistiani, Djamaluddin & Rahim, 2022). The method of using IpTT is that the tip of the patient's toe is touched with the examiner's finger. Touch for 1-2 seconds is gentle and smooth and no pushing, tapping, or piercing is given to the six points of the patient's toes who have been asked to close the eyes during the examination. Foot sensitivity assessment at 6 locations including the tip of the right big toe, the tip of the little toe of the right foot, the tip of the left big toe, the tip of the little finger of the left foot, the tip of the middle finger of the right foot, the tip of the middle finger of the left foot. If touch 2 is no sensation, there may be damage to the neurosensory peripherals. The maximum value is 6 and the minimum value is 0. The statistical test uses descriptive numerals.

RESULTS

Characteristics of Respondents in DM Type 2 Patients at the Kebonsari Health Center Surabaya

Table 1: Characteristics of Respondents in DM Type 2 Patients at the Kebonsari Health Center Surabaya in May 2020

No.	Characteristics of Respondents	Frequency	Percentage
1.	Gender	Man	18 45%
		Women	22 55%
2.	Age	49-55 years	5 12.5%
		56-61 years	16 40%
		62-67 years	19 47.5%
3.	Last education	Elementary school	6 15%
		Junior high school	12 30%
		Senior high school	20 50%
		College	2 5%
4.	Job Experiences	Farmworkers	7 17.5%
		Housewife	16 40%
		Government employees	3 7.5%
		Pensioner	1 2.5%
		Enterpriser	13 32.5%
6	Cigarette consumption	Yes	11 27.5%
		No	29 72.5%
7	Diabetic Foot Gymnastics	Yes	10 25%
		No	30 75%
8	Take DM drugs	Regular	36 90%
		Irregular	4 10%
9	Blood Sugar Control	Regular	31 77.5%
		Irregular	9 22.5%

Table 1 shows that the characteristics of respondents in DM type 2 sufferers are mostly female, aged 62-67

years, last high school education, work as a housewife, do not consume cigarettes, do not exercise diabetic feet, take DM medication, and control blood sugar regularly.

Sensory nerve examination results with IpTT examination in DM type 2 patients at the Kebonsari Health Center Surabaya

Table 2: Sensory Nerve Examination Results with IpTT Examination in DM Type 2 Patients at the Kebonsari Health Center Surabaya In May 2020

IPTT Score (There is a sensation)	Frequency	Percentage
0	0	0%
1	0	0%
2	3	7.5%
3	14	35.0%
4	12	30.0%
5	8	20%
6	3	7.5%
Total	40	100%
Average value	3.85	

Table 2 shows that the 40 respondents who underwent neurosensory peripheral examination using IpTT had a score of 2 as many as 3 people (7.5%), a score of 3 as many as 14 people (35.0%), a score of 4 as many as 12 people (30.0%), a score of 4 5 as many as 8 people (20.0%) and score 6 as many as 3 people (7.5%). The average value of the neurosensory peripheral examination using IpTT was 3.85.

DISCUSSION

Based on the research data obtained in table 2, in general, the results of the neurosensory peripheral examination in patients with diabetes mellitus type 2 through the IpTT examination have an average value of 3.85 out of a total score of 6, which means that the respondent has decreased sensory function. This is because people with diabetes mellitus experience sensitivity disorders due to damage to large nerve fibers. These nerve fibers supply the distal part of the foot and cause the foot to lose light and touch sensation. Increased oxidative stress will interfere with impulse delivery so people with diabetes mellitus will lose foot protection. Symptoms that arise will make people with diabetes mellitus not feel any touch or pressure on the tips of the toes (Deli *et al.* 2014).

According to Bare & Smeltzer (2002) loss of protective sensation of the feet is caused by chronic hyperglycemia which disrupts cell metabolism

resulting in microangiopathy and macroangiopathy. This disorder causes a decrease in blood flow to nerve cells and tissues, which can lead to hypoxia in nerve cells. Nerve cell hypoxia can cause myelination and axon stasis in nerve cells and Schwann cells so that nerve conduction can be disrupted (Sanjaya, Yanti & Puspita, 2019). Disturbances in the sensory nerve fibers/conductors of the feet have an impact on the decrease in the sensitivity of the nerves in the feet that function as protective sensations. Loss of protective sensation makes DM patients more prone to diabetic foot ulcers (Sanjaya, Yanti & Puspita, 2019).

The results of research conducted by Purwanti & Maghfirah (2016) using IpTT, showed as many as 29 people with diabetes mellitus with a percentage (85.3%) experiencing impaired foot sensitivity. Another study using IpTT said that 52 out of 93 with a percentage (55.9%) of diabetic ulcer patients had impaired foot sensitivity. Based on the researcher's observations in table 2, the assessment of peripheral neurosensory function using IpTT examination to assess foot sensitivity at 6-point locations (tip of the right big toe, tip of the little finger of the right foot, tip of the left big toe, tip of the little finger of the left foot, tip of the middle toe of the foot right, middle fingertip of left foot) the majority had decreased neurosensory peripheral function. As evidenced by the results of the IpTT score obtained a score of 2 to 4 as many as 26 respondents from 40 respondents. This means that respondents cannot feel two or more touches on their toes, then the patient is said to have decreased sensation and is at risk of developing diabetic foot ulcers.

Based on the researcher's observations in table 1, the decrease in neurosensory peripheral function is also influenced by several factors, including gender, and age, and the majority of respondents do not exercise diabetic feet. Data on the characteristics of the respondents in table 1, the female gender tends to be more, as many as 22 people (55.0%) and as many as 18 people (45.0%) are males. This is because women tend to be more at risk of developing diabetes mellitus associated with a large body mass index and menstrual cycle syndrome and during menopause which results in easy accumulation of fat which results in inhibition of glucose transport into cells (Imelda, 2019).

Age can also affect a person's sensory level decline. Based on table 1, most of the respondents aged 62-67 years were 19 people (47.5%), while those aged 56-61

years were 16 people (40.0%), while those aged 49-55 years were 5 people (12.5%). The results of this study prove that with increasing age, the risk of neuropathic complications increases (Putri & Waluyo, 2019). This is because insulin resistance increases with advancing age. Factors that contribute to insulin resistance are decreased muscle mass (sarcopenia), being overweight, reduced physical activity, and the pancreas not functioning as well as in younger people (Sujana, 2019). It is different from Kamaru, Mohd & Hussein (2018), research showed that there was no significant relationship between age, duration of diabetes, body image, occupation, monthly income, and level of education with the knowledge and the level of foot care.

Based on the characteristics of the respondents in table 1, the majority of respondents did not do diabetic foot exercises as many as 30 people (75%), and those who did diabetic foot exercises as many as 10 people (25%). Respondents who do not do diabetic foot exercises are at risk of decreased peripheral neurosensory function because they do not move, the sensation in the nerves of the feet will die, but if they do movement or exercise then there is stimulation of peripheral blood flow to increase so as not to worsen the level of neuropathy to decrease. Diabetic foot exercises can increase the use of glucose in the muscles, many cell capillaries are open so that insulin receptors become more active. This can affect controlled blood glucose levels (Sembiring, Simbolon & Lase, 2018). Diabetic foot exercise helps improve blood circulation, which carries oxygen and nutrients to the cells and nerve tissue, which will affect the metabolic processes of Schwann cells so that axon function can return to normal. Optimal nerve cell function in DM patients will maintain the function of foot sensitivity (Brahmantia *et al.*, 2020). Diabetic foot exercise can be used as an alternative intervention to increase the value of the ABI (Ankle Brachial Index) in patients with type 2 DM (Bakara & Kurniyati, 2021).

Based on the data in table 2, although the majority of respondents experienced a decrease in neurosensoriperipheral function, some respondents did not experience a decrease in (normal) neurosensoriperipheral function. These respondents had an IpTT score of 3 people (7.5%) and an IpTT 5 score of 8 people (20%) from 40 respondents, which means that respondents can feel the touch of all their toes or do not feel the touch of only one of their toes, Then the patient is said to have normal foot sensation and is not at risk of developing diabetic foot ulcers.

This is because respondents do not smoke, regularly control blood sugar and regularly take DM medication as evidenced by table 1 that respondents who do not smoke are 29 people (72.5%) and respondents who smoke are 11 people (27.5%). Respondents who regularly take DM medication are 36 people (90%) and respondents who do not regularly take DM medication are 4 people (10%). Respondents who regularly control blood sugar as many as 31 people (77.5%) and who do not regularly control blood sugar were 9 people (22.5%).

Respondents who do not smoke help improve blood pressure in the ankle and have a large effect on reducing complications (Simatupang, Pandelaki & Panda, 2013). Complications that can occur are angiopathy which causes a decrease in blood flow to nerve cells and tissues that it can cause hypoxia in nerve cells. Nerve cell hypoxia can cause myelination and axon stasis in nerve cells and Schwann cells so that nerve conduction can be disrupted (Sanjaya, Yanti & Puspita, 2019). (Respondents who are obedient or regularly take DM medication will control their blood sugar, therefore, being obedient or regularly taking DM medication can control blood sugar to prevent complications (Nanda, Wiryanto & Triyono, 2018). Knowledge and adherence to medication are very important for controlling diabetes (Al-Amedy, Yoong & Ahmed Saido, 2018).

CONCLUSION

The majority of patients with diabetes mellitus type 2 at the Kebonsari Public Health Center in Surabaya have decreased neurosensory peripheral function, so it will be a high risk of diabetic foot ulcers.

Conflict of Interest

The authors declare that they have no conflict of interests.

ACKNOWLEDGEMENT

The authors are thankful to the institutional authority for completion of the work.

Ethical Approval

This current research received ethical approval from the Health Research Ethics Committee of Sekolah Tinggi Ilmu Kesehatan Hang Tuah Surabaya, Indonesia, with a Certificate of Ethical Eligibility Number:PE/35/ VI/2020/KEPK/SHT, dated 05th June 2020.

REFERENCES

- Abo deif, H. Ibrahim, & Abdelaziz, S. H. (2019). Knowledge and Practice of Foot Care in Patients with Type 1 and 2 Diabetes at National Institute of Diabetes and Endocrinology in Cairo. *The Malaysian Journal of Nursing*, 11(02), 77–86. <https://doi.org/10.31674/mjn.2019.v11i02.009>
- Al-Amedy, O. S., Yoong, T. L., & Ahmed Saido, G. (2018). Medication Knowledge and Adherence Among Type II Diabetes Mellitus Patients: A Cross Sectional Study. *The Malaysian Journal of Nursing*, 10(02), 24–33. <https://doi.org/10.31674/mjn.2018.v10i02.003>
- Alport, A. R., & Sander, H. W. (2012). Clinical approach to peripheral neuropathy: anatomic localization and diagnostic testing. *CONTINUUM: Lifelong Learning in Neurology*, 18(1), 13–38.
- Bakara, D. M., & Kurniyati, K. (2021). Effect of Leg Exercise on the Ankle Brachial Index (ABI) of Type 2 Diabetes Mellitus Patients in Rejang Lebong Regional Hospital. *The Malaysian Journal of Nursing*, 13(2), 50–55. <https://doi.org/10.31674/mjn.2021.v13i02.009>
- Bare & Smeltzer. (2002). *Buku Ajar Keperawatan Medikal Bedah Brunner & Suddart (Alih bahasa Agung Waluyo) (8th ed.)*. EGC.
- Basir, I. S., Syam, Y., Yusuf, S., & Sandi, S. (2020). Accuracy of Ipswich touch test (IpTT) to detect small fiber neuropathy and large fiber neuropathy as a risk factor of diabetic foot ulcers in public health centers. *Enfermeria Clinica*, 30, 308–312. <https://doi.org/10.1016/j.enfcli.2019.07.108>
- Brahmantia, B., Falah, M., Rosidawati, I., Sri R, A., & Dinia F, N. (2020). Pengaruh Senam Kaki Diabetik Terhadap Sensitivitas Kaki Penderita Dm Di Puskesmas Parungponteng Kecamatan Cibalong Kabupaten Tasikmalaya. *Healthcare Nursing Journal*, 2(2), 15–19. <https://doi.org/10.35568/healthcare.v2i2.862>
- Deli, G., Bosnyak, E., Pusch, G., Komoly, S., & Feher, G. (2013). Diabetic neuropathies: diagnosis and management. *Neuroendocrinology*, 98(4), 267–280. <https://doi.org/10.1159/000358728>
- Diabetes, U. K. (2012). Touch the toes test. *Diabetes UK: London*.
- Hicks, C. W., & Selvin, E. (2019). Epidemiology of peripheral neuropathy and lower extremity disease in diabetes. *Current Diabetes Reports*, 19(10), 1–8. <https://doi.org/10.1007/s11892-019-1212-8>
- Imelda, S. I. (2019). Faktor-faktor yang mempengaruhi terjadinya diabetes melitus di Puskesmas Harapan Raya tahun 2018. *Scientia Journal*, 8(1), 28–39. <https://doi.org/10.35141/scj.v8i1.406>
- Kamaru Z. N. H., Mohd S. N. S., & Hussein S. Z. (2018). Knowledge and practice of foot care among diabetic elderly in UKM Medical Centre (UKMMC). *The Malaysian Journal of Nursing*, 9(3), 6–12.
- McLaren, A. M., & Lu, S. H. (2022). Interprofessional Validation of the Ipswich Touch Test in Adults with Diabetes: The Canadian Experience. *Canadian Journal of Diabetes*. <https://doi.org/10.1016/j.jcjd.2022.07.003>
- Nanda, O. D., Wiryanto, B., & Triyono, E. A. (2018). Hubungan Kepatuhan Minum Obat Anti Diabetik dengan Regulasi Kadar Gula Darah pada Pasien Perempuan Diabetes Mellitus. *Amerta Nutrition*, 2(4), 340. <https://doi.org/10.20473/amnt.v2i4.2018.340-348>
- Purwanti, L. E., & Maghfirah, S. (2016). Faktor Risiko Komplikasi Kronis (Kaki Diabetik) dalam Diabetik Melitus Tipe 2. *The Indonesian Journal of Health Science*, 7(1), 26–39.
- Putri, R. I. (2015). Faktor Determinan nefropati diabetik pada penderita diabetes mellitus di RSUD Dr. M. Soewandhie Surabaya. *Journal Berkala Epidemiologi*, 3(1), 109–121.
- Putri, R. N., & Waluyo, A. (2020). Faktor Resiko Neuropati Perifer Diabetik Pada Pasien Diabetes Mellitus Tipe 2: Tinjauan Literatur. *Journal Keperawatan Abdurrah*, 3(2), 17–25. <https://doi.org/10.36341/jka.v3i2.839>

- Riskesdas, L. N. (2018). Kementerian Kesehatan RI Badan Penelitian dan Pengembangan Kesehatan. <https://ejournal2.litbang.kemkes.go.id/index.php/lpb/article/view/3757>
- Rosli, N. A., Hj Md Sofian, H. N., Md Saiful, F. H., & Zolkefli, Y. (2022). Nurses' Perceptions of the Responsibility of Patients with Diabetes Mellitus: a Vignette Study. *The Malaysian Journal of Nursing*, 13(03), 19–25. <https://doi.org/10.31674/mjn.2022.v13i03.003>
- Sanjaya, P. B., Yanti, N. L. P. E., & Puspita, L. M. (2019). Pengaruh Senam Kaki Diabetik Terhadap Sensitivitas Kaki Pada Pasien Dm Tipe 2. *Community of Publishing in Nursing (COPING)*, 7(2), 97-102.
- Sembiring, E. E., Simbolon, P., & Lase, E. (2018). Pengaruh Senam Kaki Terhadap Penurunan Neuropati Pada Pasien Dengan Luka Kaki Diabetik di Asri Wound Care Medan. *Journal Mutiara Ners Juli*, 1(2), 114-120.
- Sharma, S., Kerry, C., Atkins, H., & Rayman, G. (2014). The Ipswich Touch Test: a simple and novel method to screen patients with diabetes at home for increased risk of foot ulceration. *Diabetic Medicine*, 31(9), 1100-1103. <https://doi.org/10.1111/dme.12450>
- Simatupang, M., Pandelaki, K., & Panda, A. L. (2013). Hubungan Antara Penyakit Arteri Perifer Dengan Faktor Risiko Kardiovaskular Pada Pasien Dm Tipe 2. *E-Clinic*, 1(1), 7–12. <https://doi.org/10.35790/ecl.1.1.2013.1179>
- Soheilykhah, S., Rashidi, M., Dehghan, F., Shojaoddiny-Ardekani, A., & Rahimi-saghand, S. (2013). Prevalence of peripheral neuropathy in diabetic patients. *Iranian Journal of Diabetes and Obesity*, 5(3), 107-113.
- Sujana, T. (2019). Peran Puskesmas Dalam Identifikasi Dini Penyakit Diabetes Melitus Pada Lansia. *Jurnal Kesehatan Bakti Tunas Husada: Journal Ilmu-Ilmu Keperawatan, Analisis Kesehatan Dan Farmasi*, 19(1), 111–123. <https://doi.org/10.36465/jkbth.v19i1.456>
- Sulistiani, I., Djamaluddin, N., & Rahim, N. K. (2022). Skrining Kaki Diabetes “Ipswich Touch Test (IpTT)” Dalam Mendeteksi Resiko Luka Kaki Pada Pasien DM. *Borneo Community Health Service Journal*, 2(2), 51-56. <https://doi.org/10.35334/neotyce.v2i2.2815>
- Tabatabaei-Malazy, O., Mohajeri-Tehrani, M. R., Madani, S. P., Heshmat, R., & Larijani, B. (2011). The prevalence of diabetic peripheral neuropathy and related factors. *Iranian Journal of Public Health*, 40(3), 55-62.

HOME / Guide to Authors

Guide to Authors

Guidelines for Contributing Articles to The Malaysian Journal of Nursing (MJN)

The MJN is the journal reflecting views, reports, matters of interest, and updates within the various specialties in Professional Nursing – including Clinical Nursing (which itself has various sub specialties e.g., Psychiatric Nursing, Critical Care Nursing, Oncologic Nursing, etc.), Education (all aspects relevant to nursing), Management, Community Health and Research, among others. This journal welcomes **ORIGINALLY WRITTEN ARTICLES** by contributors, **CASESTUDIES**, **LITERATURE REVIEWS**, **ANY COMMENTARY RELEVANT TO PROFESSIONAL NURSING**, and **RESEARCHES/ EVIDENCES** that directly affect the professional practice in nursing-be it clinical or otherwise.

There is no article submission charges. But after selection of the article for publication the author has to pay article processing charges.

Journal Publication Charges:

There is no Publication fee for the subscribers

Subscribe with the Malaysian Journal of Nursing and get free publication:

The publication charges will be waived if any of the author of the article subscribe for the journal. The subscriber is entitled to get 4 hard copies of the journal per year. Subscription rates 50 USD/year (plus postal charges as applicable)

Your subscription includes:

1. Cross References DOI
2. Lifetime Access to the published online articles
3. Free plagiarism checking and editing

Publication Fee for Non Subscribers:

Journal Publication Charges: USD 75 *

The charges include:

- . Plagiarism checking, editing, and draft preparation along with Cross Ref DOI charges.

- . The article will be given free open access.
- . Inclusion of the paper in printed form of hard copy journal.

Author can buy their copy at the rate USD25 including postal charges.

*Publication fees waiver is available for authors of developing countries upon request.

Submitting Articles and Research Papers

1. Articles should contain approximately 1000 to 4000 words. Shorter or longer submissions may be submitted upon approval from the editor.
2. Articles must be [submitted online](mailto:info@mjn.com.my) or as an email attachment to: info@mjn.com.my.
3. Articles (including the references) should have the following specifications:
 - i. Text document must be in Microsoft Word compatible format.
 - ii. Body text must be double-spaced, in font size 12pt and Times New Roman; Paragraphs are right justified and make sure that all pages are clearly numbered.
 - iii. Title must be font size 16pt, in bold and Times New Roman.
 - iv. Subheading must be font size 14pt, in bold and Times New Roman.
4. Plagiarism- All articles will be checked for plagiarism. If plagiarism is found the article will be rejected.
5. Segment the Articles in the following parts:
 - a. Title Page :
 - i. Title of article.
 - ii. The full names of the authors.
 - iii. Job title/Occupation, employer, location and affiliation of each author.
 - iv. Name and address of the author responsible for correspondence (Include address, telephone number and email address).
 - b. Abstract: Provide a 200 to 250-word glimpse about the article's framework/outline, which will include major findings if it is a research article.
 - c. Tables, Illustrations, Figures and Photographs : Graphics such as tables, illustrations, figures and photographs that are in low resolution must be embedded in the article you are submitting (in MS word). High resolution version of the graphics must be e-mailed to us as additional attachment.
 - d. References: Use the APA Referencing Guide.
 - e. Consent: If you are publishing photographs of patients, kindly secure their consent. E-mail us a copy of the agreement along with your article or research papers.
 - f. Copyright waiver: Along with the article, the copyright waiver statement duly signed by author and co-author(s) must be submitted.

All articles published in The Malaysian Journal of Nursing will have copyrights under MJN.

[Copyright Waiver Form](#)

g. Originality and Plagiarism: The authors should ensure that they have written entirely original works, and if the authors have used the work and/or words of others, that must be appropriately cited or quoted. All articles will be checked for plagiarism. If plagiarism is found high the article will be rejected.

General Instructions for Author's:

Types of Articles:

Original Articles: Should contain title page, abstract, keywords, introduction, research methods, results and discussion, conclusion, acknowledgement, and references. The length of the text should be limited to 12-22 pages including the references.

Review Articles: Should be requested by the editor, but MJN will also accept submitted reviews. The authors of review articles are invited to contact the Editorial Office before preparing a review article. Both solicited and unsolicited review articles are subjected to editorial review such as the original papers.

Letters to the Editor: MJN accepts letters to the editor. Letters, less than 500 words, should discuss articles published in the journal during the previous six months. Letters will undergo peer-review processing and will be edited for clarity.

Case Study: It is included all the reports about the special issues and new procedure in some limited cases with good results but not confirmed yet internationally as a global choice. It should be not exceeded more than 2000 words.

Structure of Articles:

Title Page: Should contain title which should include the study design, author(s) information such as the first name, last name, highest academic degree, and affiliation.

Abstract: Should contain background, research methods, results, and conclusion sections separately. The objective of study, findings (including its statistical significance) and the conclusion made based on the findings should be clearly presented.

Abstract is not necessary for case reports, while review articles should have an abstract. Nevertheless, the length of an abstract should be less than 250 words. Authors need to be careful that the abstract reflects the content of the article accurately.

Keywords: Each submitted article should contain 3-4 keywords.

Keywords will assist indexers in cross indexing the article as they are published with abstract.

Introduction: This should summarize the purpose and the rationale for the study. It should neither review the subject extensively nor should it have data or conclusions of the study.

Research Method: This should include exact method or observation or experiment. Mathematical and Statistical methods must be mentioned and specify any general computer package used.

Results and Discussion: Results must be presented in the form of text, tables, and illustrations. The contents of the tables should not be all repeated in the text. Instead, a reference to the table number may be given. Long articles may need sub-headings (mentioned on page 1 as Subdivisions) within some sections to clarify their contents.

Discussion should emphasize the present findings and the variations or similarities with other work done in the field by other researchers. The detailed data should not be repeated discussion again. Emphasize the new and important aspects of the study. It must be mentioned whether the hypothesis mentioned in the article is true, false or no conclusions can be derived.

Conclusion: Your conclusion is your chance to have the last word on the subject. The conclusion allows you to have the final say on the issues you have raised in your paper, to summarize your thoughts, to demonstrate the importance of your ideas, and to propel your reader to a new view of the subject. It is also your opportunity to make a good final impression and to end on a positive note. In conclusion you represent something important to your readers, and can be used for some, or all the following tasks:

- Emphasizing the purpose and importance of your article
- Explaining the significance or consequences of your findings
- Indicating the wider applications of the method developed in your article
- Establishing your article as the basis for further investigation

Acknowledgement: All contributors who do not meet the criteria for authorship should be covered in the acknowledgement section. It should include persons who provided technical help, writing assistance and departmental head that only provided general support. Financial and material support should also be acknowledged.

Author(s) must acknowledge and declare any sources of funding and potential conflicting interest, such as receiving funds or fees by, holding stocks and shares in and organizations that may profit or lose through publication of your paper. Declaring a competing interest will not lead to automatic rejection of the paper, but we would like to be made aware of it.

Submission: Manuscripts must have been written in English. Also the manuscript must have confirm letter that is assumed manuscripts are exclusively submitted to MJN and have not been previously published elsewhere (except in the form of an abstract or as part of a published lecture, review or thesis) and are not under consideration by any other journal. In the covering letter one author should be specified as the “corresponding author” and all other authors should personally sign the covering letter. Authors are responsible for all statements made in their work. The editorial board has the right to insert any necessary changes so that the manuscript is harmonized with the editorial framework of the journal. All submissions to the MJN should contain a completed copy of signed covered letter containing the copyright agreement.

Final Checklist: The authors must ensure that before submitting the manuscript for publication, they have taken care of the following:

1. Title page should contain title, name of the author(s), their qualifications, mailing address for future correspondence, email address, phone, and fax number.
2. Abstract in structured format up to 250 words.
3. Keywords contain 3-5 words.
4. References should be typed as instruction that mentioned above.
5. Tables and figures should be typed in appropriate point in the article rather than the end of the article.
6. Copyright and covering Letter.

Submitting Letters and Comments

This is open access journal, you may use the guidelines in submitting articles and research papers. Address your viewpoints, and comments to the Editor of MJN in the contacts provided herein.



Indexed In





This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Print ISSN: 2231-7007 Online ISSN: 2462-246X

©2021 THE MALAYSIAN JOURNAL OF NURSING (MJN) – A Journal for the Healthcare and Nursing Community
All Rights Reserved (website design).

Platform &
workflow by
OJS / PKP

