p-ISSN : 2686-0538 e-ISSN : 2686-2123



International Journal of Nursing and Midwifery Science Volume 5 Issue 2 | August 2021



Q Search



Announcements

Current Archives

Additional Menu 🕤 🛛 About

Home / Editorial Team

Editor In Chief

Amar Akbar, S.Kep, Ns, M.Kes. Ph.D (c), Bina Sehat PPNI Institute of health Science, Indonesia

Reviewer

Professor Denise Fassett, BHS.RN.MN.PhD.Faculty Of Health, Departement Nursing and Midfiwifery, University of Tasmania, Australia

Professor John Christian Fox, MD, RDMS, FAIUM, School Of Medicine, University Of California, Irvine, US

Prof. Eileen Savage, School of Nursing and Midwifery, University College Cork, Ireland

Associate Professor Linda Sweet, School of Nursing and Midwifery Flinders University Australia, Australia

Professor Dr. Nursalam, M.Nurs (Hons), Faculty Of Nursing Airlangga University, Indonesia

Dr. Ah. Yusuf, S.Kp.M.Kes, Faculty Of Nursing Airlangga University, Indonesia

Ferry Efendi, S.Kep., Ns., MSc, PhD, Faculty Of Nursing Airlangga University, Indonesia,

Joko Gunawan, S.Kep.Ns MN, Ph.D., Chulalongkrong University, Thailand,

Dr.Indah Lestari, S.Kep.Ns.M.Kes, Bina Sehat PPNI Institute of health Science, Indonesia

Dr.Noer Saudah, S.Kep.Ns.M.Kes, Bina Sehat PPNI Institute of health Science, Indonesia

Lely Lusmilasari, S.Kp.M.Kes.Ph.D, Faculty Of Nursing Gajah Mada University, Indonesia

Dr.H. Ahsan, S.Kp.M.Kes, Faculty Of Nursing Brawijaya University, Indonesia

Dr. Ari Natalia Probandari, Faculty of Medicine, Universitas Sebelas Maret, Indonesia, Indonesia

Dr. Farhan Alshammari, College of Nursing, University of Hail, Saudi Arabia

Dr. Bahrul Ilmi, Skp, M.Kes. Goverment School Of Nursing Medan, Indonesia

Prof. Dr. Husaeni, Skp, SKM, M.Kes, Faculty Of Public Health Lambung Amangkurat, Indonesia

Rian Adi Pamungkas, S.Kep. Ns., MNS., Dr.PH, Mahidol University, Thailand

Ni gusti ayu pramita aswitami, SST., M.Keb , Midwifery Science School of health Bina Usada Bali, Indonesia

Technical Editors :

Lasiyati Yuswo Yani, Bina Sehat PPNI Institute of health Science, Indonesia

Tria Wahyuningrum, Bina Sehat PPNI Institute of health Science, Indonesia

Section Editor


H.Anwar Sadad, S.Kom, Bina Sehat PPNI Institute of health Science, Indonesia

Eko Bagus Febriari, Bina Sehat PPNI Institute of health Science, Indonesia

Proof Reading

Asef Wildan Munfadilla, M.Pd, Bina Sehat PPNI Institute of health Science, Indonesia





ADDITIONAL MENUS Home Editorial Team Peer Review Process Focus and Scope Publication Ethics Duline Submission Guidelines Licence Term Complete Issue Download Publication Fees Submissions Indexing Copyright Notice Screen for Plagiarism Digital Archiving Policy

MAKE A SUBMISSION



View My Stats

AUTHOR GUIDLINES PACK



MAIN MANUSCRIPT TEMPLATE



This work is licensed under a Creative

Commons Attribution 4.0 International License.





Information

For Readers

For Authors

For Librarians

Editorial Office :

Editorial Office : Departement Of Research and Social Engagement Bina Sehat PPNI Institute Health Of Science, Indonesia | A Building 2nd Floor | Jl. Raya Jabon Km. 06 Mojoanyar Mojokerto. Telp/Fax.0321-390 203 | | Email: Ippmstikesbinasehatppni@gmail.com | <u>Ippm.stikes-ppni.ac.id</u>

powered by OJS Open Journal Systems

urrent Archives Announcements Additional Menu - About -	Q Search
Home / Archives / Vol 5 No NO 2 (2021): VOLUME 5 ISSUE 2 AUGUST 2021	NEED HELP (CONTACT US)
IJNMS	Via Whatshop
International Journal of Naming and Midware Science	NEED HELP (CONTACT 2)
	Via WhatsRpp (
samata	NEED HELP (CONTACT 3, for Order print Version)
	Klik Untuk Chat.
Articles	ADDITIONAL MENUS
The Increasing The Introduction Of Balanced Nutritious Food Through Cooking	Editorial Team
Class Activity On Children Of 3-4 Years Old At Ppt Pelangi Kecamatan Pakal Surabaya	Peer Review Process Focus and Scope
Abdul Muhith Muhith, Maya Zainiyah 83-87	Publication Ethics
	Online Submission Guidelines Licence Term
A SUPPORT OF THE HUSBAND DURING THE PROCESS OF LABOR AND THE PUERPERIUM IN KUPANG DISTRICT	Complete Issue Download
MARETA BAKALE BAKOIL 88-101	Publication Fees Submissions
A PDF	Indexing
	Copyright Notice
THE EXCESSIVE USE OF SOCIAL MEDIA AND ITS RELATIONSHIP TO INSOMNIA AMONG NURSING STUDENTS	Screen for Plagiarism Digital Archiving Policy
deboorah Siregar, Adellia Adellia, Alserina Hideni Harefa, Yonavia Oriza, Deborah Siregar, 102-108 Elissa Hutasoit	Reviewer
A PDF	
FACTORS AFFECTING COPING STRATEGIES FOR HEALTH WORKERS IN DEALING	
STIGMA DUE TO COVID-19 Ana Zakiyah, Ika Ainur Rofi'ah 109-116	MAKE A SUBMISSION
PDF	
SELF CARE ABILITY INFLUENCES THE QUALITY OF LIFE CONGESTIVE HEART	
FAILURE (CHF) PATIENTS rizka febtrina 117-124	🔊 sînta 🛚 🖓
D PDF	Sinca
Lemon Aromatherapy as an alternative to reduce the intensity of chemotherapy-	
related nausea and vomiting experienced by the breast cancer patients Puji Hastuti, Yuli Nurhayati, Dwi Ernawati, Christina Yuliastuti, Merina Widyastuti 125-130	View My Stats
DF	
PSYCHOLOGICAL STRESS ON NURSE PERFORMANCE IN PROVIDING NURSING	AUTHOR GUIDLINES PACK
CARE DURING THE COVID-19 PANDEMIC Meirizal Ari Putra 131-138	GUIDE for
PDF	
	MAIN MANUSCRIPT TEMPLATE
THE MOZAIC EFFECT ON THE IMPROVEMENT OF FINE MOTOR SKILLS OF 3-4 YEARS OLD CHILDREN IN INTEGRATED HEALTH CARE CENTER, THE WORKING	
AREA OF PUBLIC HEALTH CENTER WAENA JAYAPURA CITY. Suryati Romauli 139-148	DOC Paper Template
De PDF	
	@ 0
THE CORRELATION OF FAMILY KNOWLEDGE AND BEHAVIOR IN CONTROLLING HYPERTENSION IN PATIENTS AT OUTPATIENT DEPARTMENT	This work is licensed under a Creativ
	This work is licensed under a Creativ Commons Attribution 4.0 Internation License.

© 2021 International Journal of Nursing and MidwiferyScience(IJNMS)

This is an Open Access article distributed under the terms of the <u>Creative Commons</u> <u>Attribution 4.0 International License</u> which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

http://ijnms.net/index.php/ijnms

e- ISSN: 2686-2123 p- ISSN: 2686-0538



ORIGINAL RESEARCH

LEMON AROMATHERAPY AS AN ALTERNATIVE TO REDUCE THE INTENSITY OF CHEMOTHERAPY-RELATED NAUSEA AND VOMITING EXPERIENCED BY THE BREAST CANCER PATIENTS

Puji Hastuti¹, Yuli Nurhayati², Dwi Ernawati³, Christina Yuliastuti⁴, Merina Widyastuti⁵ Stikes Hang Tuah Surabaya

Email : pujihastuti@stikeshangtuah-sby.ac.id, pujihastutishtsby@gmail.com

ABSTRACT	Keywords
The mechanism of breast cancer is the cells growing and breeding becomeappear abnormal tissue of breast. One of the common treatments for it is chemotherapy using cytotoxic drugs. However, chemotherapy may cause nausea and vomiting as its side effects. Lemon aromatherapy is a complementary therapy in patients with breast cancer who experience nausea or vomiting. The study's purpose was to know the effect of lemon aromatherapy on the intensity of nausea and vomiting experienced by the breast cancer patients as an effect of chemoterapy in the Chemo Center Room of RSAL Dr. Ramelan Surabaya. The study was the pre-experimental design with pre-post test without control group. There are two variables, lemon aromatherapy is independent, and the intensity of nausea and vomiting is dependent. The sampling technique was nonprobability purposive sampling, with 34 breast cancer patients taken as the sample. A questionnaire was the instrument for collecting the data. The Data collected were analyzed using the Wilcoxon Test ($\alpha = 0,05$). The study's result indicated thatthe lemon aromatherapy was effectively to decrease of the intensity of nausea and vomiting experienced by the respondents, with the value of Wilcoxon test p < 0.001. Lemon aromatherapy stimulates the raphe nucleus to produce serotonin. Which function to generate a sense of comfort and calm. For that reason, it can be used as an alternative for taking care of nausea and vomiting experienced by patients with breast cancer as the side effect of chemotherapy.	Lemon Aromatherapy, Nausea, Vomiting, Chemotherapy

INTRODUCTION

Breast cancer mechanism is the cells growing and breeding becomeappear abnormal tissue of breast (Ismonah, 2013). The medical records from the RSAL Dr. Ramelan Surabaya showed that the number of breast cancer patients undergoing chemotherapy was 599 patients in 2015, 406 patients in 2016, and 522 patients in 2017. Studies in the USA showed that 22% - 50% of the patients undergoing chemotherapyexperienced nausea and vomiting(Mulyani & Nuryani, 2013). The number of patients who underwent chemotherapy and still experienced nausea and vomiting despite being given antiemetics was 12 out of 30 patients in July 2018. Aromatherapy is a treatment applied in addition to main major treatments interventions(Subagyo, 2013). Orange oil is one of the aromatherapies that are refreshing and useful for treating nausea, flatulence, and headaches, as well as arousing appetite (Nursalam, 2017).

METHOD

This study's designwas preexperimental design with a pre-post test without control group (selfcontrol) (Jaelani, 2009). This study was conducted in the Chemo Center Room of RSAL Dr. Ramelan Surabaya from which the population was taken. The population was the breast cancer patients who experienced nausea and vomiting due to chemotherapy, around 40 people per month on average. The sample consists of parts of the population that could be used as research subjects taken through a sampling technique. Sampling is the process of selecting the portions of the population that represent the existing population (Peoples et al., 2016). The number of breast cancer patients who experienced nausea and vomiting due to chemotherapy and met the sample requirements was 34 people taken through the technique sampling of purposive.

The variables were the lemon aromatherapy as independent while the intensity of nausea and vomiting as dependent. The intensity of nausea and vomiting was indicated by using the INVR (Index of Nausea, Vomiting, and Retching) questionnaire. The INVR questionnaire consisted of 8 questions with 5 Likert scale responses (0-4) to be filled out by therespondents(Nasif et al., 2011). The severity nausea and vomiting was measured after the first 12 hours from the insertion of drugs before chemotherapy the _ administration of lemon aromatherapy – and the next 12 hours after the administration of The lemon aromatherapy. researcher explained to the respondents about how to lemon aromatherapy. use Lemon aromatherapy should be used after 12 hours from the administration of chemotherapy drugs by dripping 2-3 drops of lemon aromatherapy essential oil on tissue paper and then inhaling it 3 times of breathing for 30 minutes.

The researcher did not observe until the process of chemotherapy was completed but returned to the respondents the next day to scale the intensity of nausea and vomiting they might experience after applying the lemon aromatherapy by distributing questionnaires. The result of the demographic questionnaire and the intensity of nausea and vomiting were made in the form of percentages and narratives. Bivariate analysis was used to identify the difference in the intensity of nausea and vomiting pre and post the lemon therapy intervention (Koensoemardiyah, 2009).

The sample size in this current study was 34 respondents. In the normality test, it was found out that the data was not normally distributed. Therefore, the data analyzedby the Wilcoxon for the intervention group with a p-value = 0.001. The significant level used was 0.05 which means that if $p < \alpha = 0.05$, the hypothesis is accepted, indicating that there is an influence of lemon aromatherapy on nausea and vomiting's intensitythat reported by the patients with breast cancerwho given the chemotherapy in the Chemo Center Room, RSAL Dr. Ramelan Surabaya

RESULTS

The study was conducted in the Chemo Center Room of RSAL Dr. Ramelan Surabaya, and the data collection was carried out on January 15-18, 2019.The respondents were 34 breast cancer patients who getting chemotherapy and has nausea and vomiting during that therapy. The demographic data were obtained through a questionnaire filled out by the respondents.

Table 1: Characteristics of RespondentsBased on the Demographic Data of theBreast Cancer Patients at the ChemoCenter Room RSAL Dr. RamelanSurabaya 15-18 January 2019 (n = 34)

Demograp	Catego	Freque		
hic data	ry	ncy (f)	age (%)	
Ages	19-29	2	5,9	
	30-40	5	14,7	
	41-51	11	32,4	
	>52	16	47,1	
Marital Status	Married	29	85,3	
	single	3	8,8	
	Divorce d	2	5,9	
Number of Children	1 child	5	14,7	
	2 children	18	52,9	
	> 2 children	7	20,6	
	Do not have children	4	11,8	
Live with	Family	32	94,1	
	Relative s	1	2,9	
	Orphan age	0	0	
	Etc.	1	2,9	
Chemother apy Cycle	2	2	5,9%	
	3	5	14,7%	
	4	16	47,1%	
	5	5	14,7%	
	6	6	17,6%	

Table 2: Characteristics of Respondents
by Drug Administration to the Breast
Cancer Patients at Chemo Center Room
of RSAL Dr. Ramelan Surabaya, 15-18
January 2019 $(n = 34)$

No	Name of	Frequen	Percenta
	Drugs given	cy (f)	ge (%)
1	Brexel, Epirol	7	20,6%

2	Carboplatin,	1	2,9%
	Dexotaxel		
3	Carboplatin,	1	2,9%
	Funcopac		
4	Carboplatin,	11	32,4%
	Paclitaxel		
5	Carboplatin,	6	17,6%
	Paxomed		
6	Cyclophospam	1	2,9%
	ide		
7	Cyclophospam	1	2,9%
	ide, brexel		
8	Cyclophospam	1	2,9%
	ide,		
	dexorubicin		
9	Cyclophospam	1	2,9%
	ide, epirubicin		
10	Cyclophospam	1	2,9%
	ide, 5FU		
11	Dexotaxel,	1	2,9%
	epirubicin		
12	Dexotaxel,	1	2,9%
	dexorubicin		
13	Paclitaxel,	1	2,9%
	epirubicin		
Tot		34	100
al			

Table The effect of Lemon 3: Aromatherapy on the Intensity of Nausea and Vomiting experienced by the BreastCancer Patients as the effect of Chemotherapy in the Chemo Center Room of RSAL Dr. Ramelan Surabaya, 15-18 **January 2019** (n = 34)

January 2017 (II - 57)				
The inten	before intervention		After intervention	
sity of naus ea and vomi ting	Frequ ency (f)	Percen tage (%)	Frequ ency (f)	Percen tage (%)
No nause a and no vomit ing	0	0	1	2,9
Mild	11	32,4	30	88,2
Medi um	21	61,8	2	5,9

Heav	2	5,9	1	2,9
у				
Total	34	100	34	100
Result of Wilcoxon Test $p = 0,000$				

The result showed that the breast cancer patients before being given lemon aromatherapy was mostly on the medium scale of nausea and vomiting, amounting to 21 respondents (61.8%). The amount changed after being given lemon aromatherapy was mostly on a mild scale, amounting to 30 respondents (88.2%). This indicated that there was a decrease in the degree of nausea and vomiting from a medium-scale to a mild scale. In other words, there is an increase in the number of respondents whose intensity of nausea and vomiting decreases from the medium scale to the mild scale after the administration of lemon aromatherapy. Based on the result of the Wilcoxon test, it was found out that there was an effect of lemon aromatherapy on nausea and vomiting's intensity that reported by the patients with breast cancer who given the chemotherapy in the Chemo Center Room of RSAL Dr. Ramelan Surabaya with a p-value of 0.001 (<0.05).

DISCUSSIONS

The result supports previous studies that conducted by Widagdo. His research showing that "Lemon aromatherapy and progressive muscle relaxation affected the decrease of the intensity of nausea and vomiting during chemotherapy"(H. & K., 2007). The benefits of lemon aromatherapy include improving mood, accelerating healing of diseases, improving health and well-being of the body, mind, and spirit (14, 12).Zorba and Isdemir (2018) explained about aromatherapy that showing in their research, "The approaches of nonpharmacological are recommended for CINV's management (Chemotherapy Induced Acute Nausea and Vomiting). It is

often combined with alternative massage"(Widagdo, medicinessuch as 2014). Potter, et al. (2011) conducted research in different groups with the number of participants are 34 the cancer patients during autologous HPC infusion given citrus therapy. The intervention group has relief nausea and vomiting level (RI, 2015). Aromatherapy does not only work when there is interference but also maintain the stability or balance of the system contained in the body. Fellowes, Barnes and Wilkinson (2004) said that aromatherapy confer short term benefits on psychological wellbeing and it may also reduce of nausea and vomiting's symptom in patients who are given treatment of cancer (D et al., 2016). The result of this current study is also supports the result of the research conducted Auwaliyah showing by that the administration of lemon aromatherapy has the effect of reducing nausea and vomiting in the first trimester of pregnancy (p =0.002)(Apriyani, 2010).

The self-nursing in providing comfort to reduce or eliminate any discomfort due to the side effects of chemotherapy is by giving complementary therapy. Aromatherapy, as a part of complementary therapy, can be used to increase of the quality of cancer patient's life(Dharma, 2011). Aromatherapy is extracted from roots, flowers, leaves, and stems of plants as well as from certain trees. One of the aromatherapies that are widely used is lemon aromatherapy (Citrus Lemon) (Setyoadi & Kushariyadi, 2011). The benefits of lemon aromatherapy include improving mood, accelerating healing of diseases, and improving health and wellbeing of the body, mind, and spirit(Subagyo, 2013). It is often combined with alternative medicines. Aromatherapy does not only work when there is interference but also maintain the stability or balance of the system contained in the body. It may also

reduce of nausea and vomiting in patients who given treatment of cancer (Widagdo, 2014). Lemon aromatherapy is one of the complementary therapies of nonpharmacological treatment to breast cancer patients who experience nausea, vomiting due to chemotherapy through the olfactory process.

CONCLUSIONS

Lemon aromatherapy reduces the intensity of nausea and vomiting experience on the patient's breast cancer as thechemotherapy's effect in the Chemo Center Room of RSAL Dr. Ramelan Surabaya. The odor will then be transmitted as a message to the olfactory center located behind the nose. Neuron cells interpret the odor and deliver it to the limbic system, which is then sent to the hypothalamus. The contents of the elements of aromatherapy will physiologically correct the imbalance that may occur in the body. The calming odor stimulates an area in the brain called the raphe nucleus to produce serotonin. It functions to generate a sense of comfort and calm, which may decrease the reported nausea of and vomiting's complaint.

ACKNOWLEDGMENT

The study has been carried out because of supportedby the breast cancer's patients, The Head of RSAL dr. Ramelan Surabaya, The Head of Cemo Center Room, and all who support until the completion of this research.

REFERENCES

Apriyani, D. (2010). Pengaruh terapi musik terhadap mual muntah lambat akibat kemoterapi pada penderita kanker pada anak usia sekolah. http://lontar.ui.ac.id/file=digital/13719 9-T Dyna Apriany.pdf

D, M., Y, S. D., & U, M. M. (2016).

Pengaruh Inhalasi Aromaterapi Lemon Terhadap Morning Sickness Pada Ibu Hamil Di Wilayah Kerja Puskesmas Tulang Bawang Tahun 2016. 2(3).

- Dharma. (2011). Metodologi Penelitian Keperawatan, Panduan Melaksanakan Dan Menerapkan Hasil Penelitian. CV. Trans Info Medis.
- H., T. T., & K., R. (2007). Obat-obat penting : khasiat, penggunaan, dan efek-efek sampingnya (ke enam). PT Elex Media Komputindo.
- Jaelani. (2009). *Aroma Therapi*. Yayasan Pustaka Obor Indonesia.
- Koensoemardiyah. (2009). Aromaterapi Untuk Kesehatan, Kebugaran Dan Kecantikan. Andi.
- Mulyani, & Nuryani. (2013). *Kanker Payudara dan PMS pada Kehamilan*. Nuha Medika.
- Nasif, Junaidi, & Muchtar. (2011). Efektivitas antiemetik pada pasien yang menggunakan sitostatika pasca bedah pada berbagai jenis kanker di rumah sakit umum daerah dr. Achmad mochtar bukit tinggi. http://jstf.ffarmasi.unand.ac.id/index.p hp/jstf/article/viewFile/56/5/9
- Nursalam. (2017). Metodologi Penelitian Ilmu Keperawatan: Pendekatan Praktis (4th ed.). Salemba Medika.
- Peoples, A. R., Roscoe, J. A., Block, R. C., H., C. E., Ryan, J. L., Mustian, K. M., & Dozier, A. M. (2016). Nausea and disturbed sleep as predictors of cancer- related fatigue in breast cancer patiens: multicenter NCORP

study. Supportive Care in Cancer. https://doi.org/10.1007/s00520-016-3520-8

- RI, P. D. dan I. K. K. (2015). *Stop Kanker*. Departemen Kesehatan.
- Setyoadi, & Kushariyadi. (2011). Terapi Modalitas Keperawatan Pada Klien Psikogeriatrik. Salemba Medika.
- Subagyo, A. (2013). *Mual dan muntah akibat kemoterapi*. http://www.klikparu.com/2013/05/mu al-dan-muntah-akibatkemoterapi.html
- Widagdo, P. A. (2014). Pengaruh aroma terapi lemon dan relaksasi otot progresif terhadap penurunan intensitas mual muntah setelah kemoterapi pada pasien kanker payudara di Rumah Sakit Telogorejo Semarang. http://ejournal.stikestelogorejo.ac.id/in

dex.php/jikk/article/view/318/341



- The submission file is in OpenOffice, Microsoft Word, or RTF document file format.
- ✓ Where available, URLs for the references have been provided.
- The text is single-spaced; uses a 12-point font; employs italics, rather than underlining (except with URL addresses); and all illustrations, figures, and tables are placed within the text at the appropriate points, rather than at the end.
- ✓ The text adheres to the stylistic and bibliographic requirements outlined in the Author Guidelines.

Author Guidelines

Author Guidelines

We now differentiate between the requirements for new and revised submissions. You may choose to submit your manuscript as a single Word file to be used in the refereeing process. Only when your paper is at the revision stage, will you be requested to put your paper in to a 'correct format' for acceptance and provide the items required for the publication of your article.

TYPES OF MANUSCRIPT

Original Articles

Original Articles should report original clinical studies or research not previously published or being considered for publication elsewhere. The text should not exceed 7000 words, including list of authors and their affiliations, corresponding author, acknowledgements and figure legends, with an abstract of maximum 250 words, a list of minimum 25 references primarily from international journal indexed by Scopus or Web of Science, and maximum 5 figures/tables (see below for more details on the layout).

Systematic Reviews

Systematic Reviews are exhaustive, critical assessments of evidence from different data sources in relation to a given subject in the areas of nursing. A systematic search of the relevant data sources should be carried out and the items collected should be carefully evaluated for inclusion based on a-priori defined inclusion/exclusion criteria. A description and an analytical graphic representation of the process should be provided. The specific features of the participants' or patients' populations of the studies included in the review should be described as well as the measures of exposure and outcome with the indication of the corresponding data sources. A structured abstract is required (like for Short reviews). The text must not exceed 7000 words including the acknowledgments, with no more than 4 tables and/or figures and minimum 40 references.

Meta-analyses

Meta-analyses should follow the same guidelines as for systematic reviews. They are expected to provide exhaustive information and statistical assessment of pooled estimates of pre-defined outcomes, study heterogeneity and quality, possible publication bias, meta-regression and subgroup analyses when appropriate. Depending on the type of study, Authors are invited to submit PRISMA flow diagrams or MOOSE checklists. Both systematic reviews and meta-analyses will be dealt with ordinarily as original articles as far as the editorial process is concerned.

Letter to Editor, Opinion, General review can use free format.

TITLE AND AUTHORSHIP


The title should describe the summary of the research (concise, informative, no abbreviation, and maximum 12 words).

Authorship of articles should be limited to those who have contributed sufficiently to take public responsibility for the contents. Full names of authors (without academic title); author's affiliation [name(s) of department(s) and institution(s)]; corresponding author's name, mailing address, telephone and fax numbers, and e-mail address. Corresponding author is the person responsible for correspondence during the publication process and post publication.

ABSTRACT

Abstracts should be less than 250 words, and should not include references or abbreviations. They should be concise and precise with enough information, highlighting the points and importance of the article. In general they should include the following:

- Introduction: One or two sentences of background and purpose of study.
- Method: describe the research design, settings (please do not mention the actual location, but use geographic type or number if necessary); Participants (details of how selected, inclusion and exclusion criteria, numbers entering and leaving the study, relevant clinical and demographic characteristics).
- Results: report main outcome(s) /findings including (where relevant) levels of statistical significance and confidence intervals.
- Conclusions: should relate to study aims and hypotheses.
- Keyword: Provide between three and five key words in alphabetical order, which accurately identify the paper's subject, purpose, method and focus.



IJNMS Has Been Indexed In :



The text should be structured as introduction, methods, results, discussion and conclusion. Footnotes are not advisable; their contents should rather be incorporated into the text. Use only standard abbreviations; use of nonstandard abbreviations can be confusing to readers. Avoid abbreviations in the title of the manuscript. The spelled-out abbreviation followed by the abbreviation in parenthesis should be used on first mention unless the abbreviation is a standard unit of measurement. If a sentence begins with a number, it should be spelled out.

ACKNOWLEDGMENT (OPTIONAL)

Acknowledgments should be limited to appropriate professionals who contributed to the paper, including technical help and financial or material support, also general support by a department chair-person.

TABLES AND FIGURES

Tables should be numbered in arabic numerals, captions should be brief, clearly indicating the purpose or content of each table. If your manuscript includes more than 5 tables in total, or for very large tables, these can be submitted as Supplementary Data and will be included as such in the online version of your article.

REPORTING GUIDELINE

Reporting guidelines endorsed by the journal are listed below:

- Observational cohort, case control and cross sectional studies STROBE Strengthening the Reporting of Observational Studies in Epidemiology, <u>http://www.equator-network.org/reporting-guidelines/strobe/</u>
 Qualitative studies - COREQ - Consolidated criteria for reporting qualitative research, <u>http://www.equator-</u>
- Qualitative studies COREQ Consolidated criteria for reporting qualitative research, <u>http://www.equator-network.org/reporting-guidelines/coreq</u>
- Quasi-experimental/non-randomised evaluations TREND Transparent Reporting of Evaluations with Non-randomized
 Designs, <u>http://www.cdc.gov/trendstatement/</u>
- Randomised (and quasi-randomised) controlled trial CONSORT Consolidated Standards of Reporting Trials, http://www.equator-network.org/reporting-quidelines/consort/
- Study of Diagnostic accuracy/assessment scale STARD Standards for the Reporting of Diagnostic Accuracy Studies, http://www.equator-network.org/reporting-guidelines/stard/
- http://www.equator-network.org/reporting-guidelines/stard/ • Systematic Review of Controlled Trials - PRISMA - Preferred Reporting Items for Systematic Reviews and Meta-Analyses, http://www.equator-network.org/reporting-guidelines/prisma/
- http://www.equator-network.org/reporting-guidelines/prisma/ • Systematic Review of Observational Studies - MOOSE - Meta-analysis of Observational Studies in Epidemiology, http://www.ncbi.nlm.nih.gov/pubmed/10789670

You can download these guidelines Author Pack Guidlines >> \underline{HERE} (.pdf).


Submission Preparation Checklist

As part of the submission process, authors are required to check off their submission's compliance with all of the following items, and submissions may be returned to authors that do not adhere to these guidelines.

- As part of the submission process, authors are required to check off their submission's compliance with all of the following items. Submissions may be returned to authors that do not adhere to these guidelines.
- The author(s) haven〙t suggested any personal information that may make the identity of the patient recognizable in any form of description, photograph or pedigree. When the photographs of the patient are essential and indispensable as scientific information, the author(s) must have received the consent in written form and have clearly stated such.
 In case of experimenting on humans, the author(s) have certified that the process of the research is in accordance with
- 3. In case of experimenting on humans, the author(s) have certified that the process of the research is in accordance with the ethical standards of the Helsinki declaration, as well as any domestic and foreign committees that preside over the experiment. If any doubts are raised as to whether the research proceeded in accordance with the Helsinki declaration, then author(s) are required to explain it. In the case of experimenting on animals, the author(s) have certified that the author(s) had followed the necessary domestic and foreign guidelines related to experimenting on animals in a laboratory.
- 4. The author(s) have received consent from the author or editor of any pictures or table quoted from other journals or books. A portion or the entirety of the article must not have been published in other journals, nor must it have contributed to other journals or is still under review.
- 5. The author(s) undersigned hereby give IJNMS the rights as publisher, the right of first publication of all published material. It will be llicensed under a Creative Commons Attribution 4.0 International License.
- It will be licensed under a <u>Creative Commons Attribution 4.0 International License</u>. 6. Author(s) of the journal have clarified everything that may arise such as work, research expenses, consultant expenses, and intellectual property on the document in relation to the <u>ICMDE form disclosure of conflicts of interest</u>.

Privacy Statement

The names and email addresses entered in this journal site will be used exclusively for the stated purposes of this journal and will not be made available for any other purpose or to any other party.



Editorial Office :

Departement Of Research and Social Engagement Bina Sehat PPNI Institute Health Of Science, Indonesia | A Building 2nd Floor | Jl. Raya Jabon Km. 06 Mojoanyar Mojokerto. Telp/Fax.0321-390 203 | | Email: lppmstikesbinasehatppni@gmail.com | |ppm.stikes-ppni.ac.id