

MALAYSIAN JOURNAL OF
Medicine and Health Sciences
Vol. 18 SUPP2 Jan 2022
Supplementary Issue:
Healthcare Challenges and
Innovations in Present Pandemic Era

Malaysian Journal of Medicine and Health Sciences Vol. 18 SUPP2, Jan 2022

MALAYSIAN JOURNAL OF

Medicine and Health Sciences

Vol. 18 SUPP2 / Jan 2022

*Supplementary Issue:
Healthcare Challenges and
Innovations in Present
Pandemic Era*



Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
43400 UPM Serdang
Selangor Darul Ehsan
Malaysia
<http://medic.upm.edu.my/>

UPM Press
Universiti Putra Malaysia
43400 UPM Serdang
Selangor Darul Ehsan
Malaysia
<http://penerbit.upm.edu.my>

PENERBIT
UPM
UNIVERSITI PUTRA MALAYSIA
P R E S S



A scientific journal published by Universiti Putra Malaysia Press

About the Journal

The Malaysian Journal of Medicine and Health Sciences (MJMHS) is published by the Faculty of Medicine and Health Sciences, Universiti Putra Malaysia. The main aim of the MJMHS is to be a premier journal on all aspects of medicine and health sciences in Malaysia and internationally. The focus of the MJMHS will be on results of original scientific research and development, emerging issues and policy analyses pertaining to medical, biomedical and clinical sciences. The Malaysian Journal of Medicine and Health Sciences is now indexed in the following data bases: Scopus, EBSCOhost, ISC, and Rubriq.

Editorial Board

Editor in Chief:

Prof. Dr. Normala Ibrahim

Members:

Prof. Dr. Rukman Awang Hamat

Prof. Dr. Christopher Lim Thiam Seong

Prof. Dr. Ching Siew Mooi

Prof. Dr. Sharmili Vidyadaran

Assoc. Prof. Dr. Rajesh Ramasamy

Assoc. Prof. Dr. Sethu Thakachy Subha

Assoc. Prof. Dr. Loh Su Peng

Assoc. Prof. Dr. Nor Afiah Mohd Zulkefli

Assoc. Prof. Dr. Wan Aliaa Wan Sulaiman

Assoc. Prof. Dr. Ling King Hwa

Assoc. Prof. Dr. Habibah Abdul Hamid

Assoc. Prof. Dr. Zubaidah Nor Hanipah

Assoc. Prof. Dr. Intan Hakimah Ismail

Assoc. Prof. Dr. Ho Yu Bin

Assoc. Prof. Dr. Karmegam Karuppiah

Assoc. Prof. Dr. Subapriya Suppiah

Lead Guest Editor:

Prof. Dr Sandeep Poddar

Guest Editors:

Prof. Dr Debajyoti Bose,

Prof. Dr. Sudipta Poddar

Prof. Dr. Krishnajyoti Goswami

Technical Coordinator:

Dr. Tan Kit Aun

International Advisory Board

Prof. Dr. Pierce Anthony Grace

Prof. Dr. David Isaacs

Prof. Dr. P. Ronan O'Connell

Prof. Dr. Graham Alexander Mc Gregor

Prof. Dr. Tan Ser Kiat

Prof. Dr. Gregory Y.H. Lip

Prof. Dr. Roger Pepperell

*Malaysian Journal of
Medicine and Health Sciences
Vol. 18 SUPP2 January 2022
Contents*

EDITORIAL

Healthcare Challenges and Innovations in COVID 19 Pandemic Era 1
Sandeep Poddar; Debajyoti Bose; Krishnajyoti Goswami; Sudipta Poddar

ORIGINAL ARTICLES

Preparation of Black Cumin Extract Nanoemulsion Using the Oil Phase of Virgin Coconut Oil (VCO) Tween 80 and PEG 400 Surfactants 2-5
Henny Syapitri, Santhna Letchmi Panduragan, Sathish Babu, Vivi Purwandari, Cut Masyitah Thaib

The Effects of Green Beans and Soybean Juice on Haemoglobin (Hb) among Female Adolescents Aged 12-14 Years 6-9
Nancy Oliy, Herinawati Herinawati, Lia Artika Sari, Enny Susilawati, Gustina Gustina, Hidayat Arifin, Sandeep Poddar

Reduction in Transaminase Enzymes Action after Application of Leaves Extract of IR Bagendit Paddy on Liver Cell of Lead Exposed Rat 10-14
Herlisa Anggraini, Budi Santosa, Erick Erianto Arif, Sandeep Poddar

Family Assistance Significantly Increased the Growth of Infants With Low Birth Weight (LBW) in the Province of Nusa Tenggara Timur, Indonesia 15-20
Florentianus Tat, Yulianti Kristen Banhae, Norma Tiku Kambuno, Faridah Binti Mohd Said, Bibi Florina Abdullah, Hafizah Che Hassan

Trends of Malaria Cases From 2018 to 2019 in East-Sumba, East Nusa Tenggara, Indonesia 21-25
Pius Selasa, Heru Santoso Wahito Nugroho, Margaretha Teli, Israfil, Saidah Rauf, Sandeep Poddar

Health Education in Reducing Community Stigma among Persons with Mental Disorders 26-30
Eni Hidayati, Enwar Usti Sumadi, Desi Ariyana Rahayu, Tri Nurhidayati, Zaidah Ismail

Walking as Simple Exercise to Reduce Blood Glucose Level in Type 2 Diabetes 31-36
Christina Yuliasuti, Setiadi, Nur Muji Astuti, Nisha Dharmayanti Rinarto

Characteristics of Multidrug-Resistant Tuberculosis and Smear-Positive Pulmonary Tuberculosis: A Case-Referent Study 37-41
Desy Indra Yani, Neti Juniarti, Mamat Lukman

Factors Correlating to Family Participation in Children Growth and Development in West Java Indonesia : A Retrospective Cohort Study 42-48
Neti Juniarti, Mamat Lukman, Citra Windani Mambang Sari, Rindang Ekawati, Dadang Suhenda, Anindita Dyah Sekarpuri

Interleukin 6 Associated with Adrenal Insufficiency in COVID-19 Patient 49-52
Nurhasan Agung Prabowo, Rohmaningtyas Hidayah Setyaningrum, Hendrastutik Apriningsih, Hartono

Emergency Room Nurse Scheduling at a Public Hospital in Jakarta during Covid-19 Pandemic, Using TPB Algorithm Method 53-58
Kosasih Bai Asmita, Santhna Letchmi Panduragan, FatimahYahya, Bibi Florina Abdullah

Dimensions of Quality of Life among Children with Congenital Heart Disease: A Survey 59-63
Ikeu Nurhidayah, Toayah Indah Sari, Aan Nuraeni

The Effect of Knowledge on Attitudes and Behavior of Nursing Students About Multivitamin Consumption During the Covid-19 Pandemic <i>Zahrah Maulidia Septimar, Jaenudin Saputra, Febi Ratnasari, Faridah Binti Mohd Said, Bibi Florina Abdullah</i>	64-66
Knowledge Survey of Coronavirus Disease (Covid-19) Prevention and Vaccination in Sukabumi, West Java, Indonesia <i>Hendri Hadiyanto, Asep Suryadin, Egi Mulyadi, Anggun Fajar Rhamadani, Mustopa Saepul, Faridah Binti Mohd Said, Hafizah Che Hassan</i>	67-69
Evaluation of Diabetes Patients About Chronic Disease Management Program in Bandung <i>Citra Windani Mambang Sari, Witdiawati Witdiawati, Dadang Purnama, Titis Kurniawan</i>	70-75
Determinants of the Final Decision to Take Children under 5 Years Old for Medical Treatment in Indonesia <i>Yoyok Bekti Prasetyo, Yulis Setiya Dewi, Hidayat Arifin, Anggraini Dwi Kurnia, Nur Lailatul Masruroh, Nur Melizza, Sandeep Poddar</i>	76-82
Relationship Between Coping Mechanisms and Quality of Life of Patients with Chronic Kidney Failure Undergoing Hemodialysis in a Hospital in the Rural Area <i>Benny Arief Sulistyanto, Irnawati, Fatma Umriana, Imron Rosyidi</i>	83-87
Effect of Health Education Through Social Media at the Level of Anxiety Pregnant Women in the Time of Covid-19 <i>Murbiah, Ika Guslanda Bustam, Rahmi Garmini</i>	88-91
Development Strategy for Pharmacy Installation at a Hospital in Bitung City Using Strengths, Weaknesses, Opportunities, and Threats (SWOT) Analysis Method <i>Gayatri Citraningtyas¹, Widya Astuty Lolo, Imam Jayanto, Anita Pratasik, Heri Wijaya, Mohammad Gousuddin</i>	92-97
Psychoeducation and Relaxation Decrease Stress Levels in COVID-19 Patients <i>Rohmaningtyas Hidayah, Nurhasan Agung Prabowo, Hendrastutik Apriningsih, Hartono</i>	98-103
Effectiveness of Occupational Therapy on Improving Eating Independence in School Age Children With Down Syndrome at Kendari Autism Service Center, Indonesia <i>Sri Wahyuni, Anisa Purnamasari, Faridah Mohd Said, Nisha Nambiar</i>	104-107
Effect of Swaddle and Conventional Tub Bath on Physiological and Comfort Response in Premature Newborns at a Government Hospital in West Java, Indonesia <i>Dior Manta Tambunan, Henny Suzana Mediani, Nunung Nurjanah, Tukimin Bin Sansuwito, Munirah Mohd Hassan</i>	108-116
Relationship Stress Levels and Parenting Mothers of Children with Autism <i>Diyah Arini, Qori'ila Saidah, Netta Meridianti Putri, Julian Dewi Praspawati, Faridah, Dwi Ernawati</i>	117-124
Effects of Cognitive Behaviour Therapy on Anxiety and Depression Reduction among Women with Surgical Menopause: A randomized Controlled Trial <i>Smitha Poovathinkal Madhavan, Indira Armugham, Amiya Bhaumik, Aruna Gundluru, Usha Kiran Thirunavukarasu, Siva Kumar A.V, Regidor III Dioso</i>	125-130
Aspects of Spirituality in Midwifery services at Az Zahra Clinic Tangerang Primary Health Services <i>Marthia Ikhlasih, Tahir Abdullah, Andi Zulkifli, Suriah, Yudhia Fratidhina, Nurhaedar Jafar, Ummu Salmah, Ridwan Amiruddin</i>	131-135
Predictive Factors of Covid-19 Infection in Resident Doctors at Specialized Isolation Center of Covid-19 Hospital <i>Ida Ayu Sri Wijayanti, Putu Gede Sudira, Ni Nyoman A. Widnyani, Putu Agus Prawira Eka Putra, Dewa Putu Gde Purwa Samatra, Komang Ayu Witarini</i>	136-141

Evaluation of Family Centered Indonesia Health Program by Using the Kirkpatrick Model on Families in South Sempaja Region Samarinda of East Borneo Province of Indonesia <i>Herni Johan Melipu, Tukimin Bin Sansuwito, Kosheila Ramuni</i>	142-146
Study of Expectant Fathers' Prejudices and Practices vs. Pregnant Women's Pleasure related to Antenatal Care at OPD, NMCH, Nellore, A.P., India <i>Aruna Gundluru, Indira Armugham, Amiya Bhaumik, Smitha Poovathinkal Madhavan, Usha Kiran thirunavukarasu, Viji Alex, Suganthi Lokonathan</i>	147-151
Health Related Quality of Life (HRQoL) as Impinge of Functional Mobility Among Elderly Living in Rural and Urban Areas, Nellore, AP - Comparative Study <i>Viji Alex, Indira Armugham, Amiya Bhaumik, V. Jayanthi, Smitha Poovathinkal Madhavan, Aruna Gundluru, Sandeep Poddar</i>	152-156
Assess 3 I's (Instruction, Initiation, Improvement) as a Road Map to Breastfeeding Success among Postnatal Mothers at NMCH, Nellore, A.P, India <i>Usha KiranThirunavukarasu, Indira Armugham, Amiya Bhaumik, Smitha P M, Aruna Gundluru, Viji Alex, Bibi Florina Abdullah</i>	157-161
Analysis of Consumer Response to Marketing Mix Strategy at Several Pharmacies in Samarinda City, East Borneo <i>Heri Wijaya, Rusdiati Helmidanora, Siti Jubaidah, Sreemoy Kanti Das</i>	162-167
Family Healthcare Tasks to Prevent the Transmission of Pulmonary Tuberculosis (TB) Among Family Having a Household Contact in Eastern Indonesia <i>Margareta Teli, Pius Selasa, Israfil, Sandeep Poddar</i>	168-175
Integrated Antenatal Care For Maternal Complications Management In Central Lampung Indonesia <i>Lusia Asih Wulandari, Zulkarnain A.Hatta, Hafizah Che Hassan</i>	176-180
Factors of Mother's Success in Exclusive Breastfeeding <i>Amellia Mardhika, Agus Sulistyono, Arina Qona'ah, Iswatun Iswatun, Joko Susanto, Anastasia Pangestu Mei Tyas</i>	181-187
Development of a Survey Instrument Mobile App to Determine the Map Location of a Drugstore <i>Irnawati, Feri Febria Laksana, Istianah, Tukimin bin Sansuwito</i>	188-194
Effect of Nurse Led Bundle Care Therapy on Pop Symptoms and Quality of Life Among Women with Prolapsed Uterus <i>JayanthiV., K. Jayashree, A. Indira, A.Viji, Rajeswari.H, Sandeep Poddar</i>	195-200
Risk Factors of Preeclampsia among Pregnant Women in Rural Area of Indonesia <i>Sri Hayati, Delia Wati Putri, Maidartati, Erna Irawan, Ramesh Prasath Rai, Ruma Poddar</i>	201-205
Stigma and Anxiety Levels With Adherence on the Treatment Schedule Patient With HIV/AIDS in Indonesia <i>Nuh Huda, Nursalam, Tintin Sukartini, Erna Heny Trisusanti, Ninik Ambar Sari, Ceria Nurhayati, Sri Anik Rustini</i>	206-212
Service Management and Cancer Patient Satisfaction in Hospitals: Cross Sectional Descriptive Study in Indonesia <i>Gina Meirawaty, Laili Rahayuwati, Kurniawan Yudianto, Habsyah Sapidah Agustina</i>	213-217
Depression, Anxiety, and Stress Levels Among Nurses in Universitas Sebelas Maret Hospital After A-Year of COVID-19 Pandemic <i>Risalina Myrtha, Nurhasan Agung Prabowo, Zahrina Mardhiyah, Aprilia Rizky Arifiani, Hendrastutik Apriningsih, Fadjri Kirana Anggarani, Hartono</i>	218-222
Adipose Tissue: All Parts Is Useful <i>In Vitro</i> Study About Adipose-Derived Mesenchymal Stem Cell <i>Budi Baktijasa Dharmadjati, Djanggan Sargowo, Aulanni'am Aulanni'am, Budi Susetyo Pikir, Oryza</i>	223-228

Sativa, Kandita Arjani, Muhammad Rifqi Djamal Hasan

The Effect of Leadership Styles on Nurse Performances and Job Satisfaction Among Nurses in Dumai Public Hospital: Technological Innovation as Mediator
Nita Ariani, Tukimin bin Sansuwito, Rames Prasath, Milya Novera, Desi Sarli, Sandeep Poddar 229-234

Dim Light Effect on Active Phase Acceleration During Labour
Rulfia Desi Maria, Risqi Utami, Reysha Melati, Lady Wizia, Evi Susanti, Tukimin Bin Sansuwito, Sandeep Poddar 235-238

The Effect of EFT (Emotional Freedom Technique) to the Self Esteem among Nurses
Nenden Lesmana Wati, Tukimin Bin Sansuwito, Ramesh Prasath Rai, Irma Darmawati, Reni Anggareni, Mayasyanti Dewi Amir, Titin Nasiatin 239-242

Occupational Hazards, the Use of Ppe, and Health Impacts Among Welders in Sumedang, West Java, Indonesia
Kosasih Bai Asmita, Santhna Letchmi Panduragan, Nisha Nambiar, Fatimah Yahya 243-244

Quality of Life Among Type Two Diabetes Mellitus Clients in an Urban Area of Tirupati
Suleman Shareef Mahammad, Armugham Indira, Amiya Bhaumik, Smitha P.M., T Usha Kiran, P Navya Keerthana, Ruma Poddar, Bibi Florina Binti Abdullah, Regidor III Dioso 245-255

Quality of Life and Its Determinants of Demographic Characteristics Among Chronic Renal Failure Patients Who Underwent Therapy of Hemodialysis in Arifin Ahmad Public Hospital, in Pekanbaru Riau Indonesia
Sofiana Nurchayati, Tukimin bin Sansuwito, Hafizah Che Hasan 256-261

Effect of Vasectomy on Quality of Life Among Adult Men in Central Lampung Indonesia
Sri Nowo Retno, Hari Krishnan Andi, Hafizah Che Hassan 262-266

The Correlation Knowledge, Attitude, and Compliance of Infection Control Guidelines at Dental Hospital Hasanuddin University
Fuad Husain Akbar, Serliawati 267-272

SYSTEMATIC REVIEW

Interventional Study on Improvement of Mothers Knowledge Regarding Danger Signs In Children Under-Five Years of Age: A Systematic Review of Literature
Baisakhi Sarkar, Anima Haldar, Nirmal Kumar Mandal, Sandeep Poddar 273-275

Comparison of Zinc Chitosan and Silver Dressing in Healing of Diabetic Ulcer: Systematic Review
Sulastri Galumbang Panahatan Tambunan, Dudut Tanjung, Imam Budi Putra, Jansen Parlaungan 276-283

Predicting SARS-Cov-2 - Procoagulant/Coagulation Related Protein Interaction: A Systematic Review
Tonang Dwi Ardyanto, Rasmaya Niruri, Adi Yugatama, Artini Pangastuti 284-290

REVIEW ARTICLE

Effectiveness of Releasing the Umbilical Cord Using Dry Gauze and the Open Technique: A Review
Febi Ratnasari, Zahrah Maulidina Septimar, AYG.Wibisono, Farzana Yasmin, Faridah Binti Mohd Said 291-294

Quality of life, Perceived Stress, Coping and Burden among caregivers of Behavioral Disorder Children: A Narrative Review
Anita Paul Samanta, Sasmita Das, Saswati Nath 295-301

Health Literacy of ICU Patients' Family: A Review
Ninuk Dian Kurniawati, Yulis Setiya Dewi, Erna Dwi Wahyuni, Hidayat Arifin, Rifky Octavia Pradipta, Sandeep Poddar 301-305

CASE REPORT

Survival of Outpatient Department Services During COVID-19 Pandemic: A Case Study
Nurhasan Agung Prabowo, Risalina Myrtha, Hendrastutik Apriningsih, Maya Permata Sari, Hartono 307-309

Lupus Flares in COVID-19 Patients: A Case Report
Nurhasan Agung Prabowo, Risalina Myrtha, Hendrastutik Apriningsih, Hartono, Gladys Sudyanto 310-312



Source details

Malaysian Journal of Medicine and Health Sciences

Scopus coverage years: from 2007 to 2023

Publisher: Faculty of Medicine and Health Sciences, University Putra Malaysia

ISSN: 1675-8544

Subject area: Medicine: General Medicine

Source type: Journal

CiteScore 2022

0.4



SJR 2022

0.139



SNIP 2022

0.197



[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology



CiteScore 2022 counts the citations received in 2019-2022 to articles, reviews, conference papers, book chapters and data papers published in 2019-2022, and divides this by the number of publications published in 2019-2022. [Learn more >](#)

CiteScore 2022 ▾

$$0.4 = \frac{758 \text{ Citations 2019 - 2022}}{1,963 \text{ Documents 2019 - 2022}}$$

Calculated on 05 May, 2023

CiteScoreTracker 2023 ⓘ

$$0.5 = \frac{1,090 \text{ Citations to date}}{2,412 \text{ Documents to date}}$$

Last updated on 05 February, 2024 • Updated monthly

CiteScore rank 2022 ⓘ

Category	Rank	Percentile
Medicine		
General Medicine	#688/830	17th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

All content on this site: Copyright © 2024 Elsevier B.V. ↗, its licensors, and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the Creative Commons licensing terms apply.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.



ORIGINAL ARTICLE

Walking as Simple Exercise to Reduce Blood Glucose Level in Type 2 Diabetes

Christina Yuliasuti, Setiadi, Nur Muji Astuti, **Nisha Dharmayanti Rinarto**

Sekolah Tinggi Ilmu Kesehatan Hang Tuah Surabaya, Jl. Gadung No.1 Surabaya 60244, Indonesia

ABSTRACT

Introduction: Patients with Type 2 Diabetes Mellitus (DM) who do not do physical exercise are more likely to be unable to control their blood glucose levels, which has the potential to cause various complications. Based on basic health research 2018, the prevalence of DM is mostly at the age of 55-64 years (15.6%). This current research aims to identify the effectiveness of walking to reduce blood glucose levels. **Methods:** This study, which applied the time-series design, involved 15 randomly selected patients with type 2 Diabetes Mellitus aged 45-65 years. The walking intervention was carried out three times in 2 weeks, namely twice in week 1 and once in week 2, each for 30 minutes. Data were collected using a glucometer and an observation sheet. Data were analyzed using paired t-test and repeated ANOVA with a significance level of $p < 0.05$. **Results:** We found differences in blood glucose levels before and after walking on days 1, 3, and 8 ($p < 0.001$). In addition, there were significant differences in all measurements of blood glucose levels of Diabetes Mellitus patients both at the start (before walking) and after walking on day 3 and day 8 ($p < 0.025$). This study found that walking was effective at lowering blood glucose levels. **Conclusion:** Walking as a simple exercise can increase the use of glucose by active muscles, reduce insulin resistance, and increase blood vessel flow. Health education on self-care management, especially physical exercise on walking, will support the control of blood glucose levels.

Keywords: Blood glucose, Diabetes mellitus, Exercise, Walking

Corresponding Author:

Christina Yuliasuti, M.Kep.

Email: christinayuliasuti@stikeshangtuah-sby.ac.id

Tel: +62 817328684

INTRODUCTION

Based on American Diabetes Association, 2005, Diabetes Mellitus (DM) is a group of metabolic diseases with the characteristic of hyperglycemia that occurs due to abnormalities of insulin secretion, insulin work, or both (1). Based on Perkeni's recommendation (Indonesian Endocrinology Association), to prevent chronic complications, effective DM control is needed. DM control applies the 4 pillars of diabetes mellitus control guidelines that involve education, dietary regulation, exercise, and medication adherence (2).

Based on the phenomenon that occurred at Puskesmas (Local Clinic) of Kebonsari in Surabaya, people with diabetes mellitus tend to be less familiar with the 4 pillars of DM control. They did not fully understand that walking is an easy and cheap way to control blood glucose levels. Several factors were inhibiting these activities, such as the patients were too busy, the patients' ignorance of the management of DM, economic reasons, and others. Based on the researchers'

knowledge, so far the effectiveness of walking to reduce blood glucose levels in diabetes mellitus sufferers in the respective research site cannot be explained.

"The number of people with diabetes rose from 108 million in 1980 to 422 million in 2014. Prevalence has been rising more rapidly in low- and middle-income countries than in high-income countries. Diabetes is a major cause of blindness, kidney failure, heart attacks, stroke and lower limb amputation. Between 2000 and 2016, there was a 5% increase in premature mortality from diabetes. In 2019, an estimated 1.5 million deaths were directly caused by diabetes. Another 2.2 million deaths were attributable to high blood glucose in 2012" (3 p. 1).

"The International Diabetes Federation (IDF) Atlas 2017 reported that the Diabetes epidemic in Indonesia showed an increasing trend. Indonesia was the sixth country in the world after China, India, the United States, Brazil, and Mexico whose around 10.3 million people aged 20-79 years suffered from Diabetes" (4 p. 1).

Based on the 2018 Basic Health Research, the prevalence of diabetes mellitus in East Java in the population of all ages was 2.0%. Meanwhile, the prevalence of Diabetes Mellitus in people aged >15 years was 2.6%.

The prevalence of diabetes mellitus based on the examination of blood glucose levels in people aged >15 years, according to the highest characteristics, is at the age of 55-64 years (15.6%) (5).

The total population in the Kebonsari Surabaya is 48,962 people. Out of which 3,360 people in 2018 were affected by Diabetes mellitus (DM). Based on the registration data in 2019, the visits of DM sufferers at Puskesmas Kebonsari in the last 3 months (January to March 2019) were around 460 people. In a preliminary study conducted by researchers on April 1, 2019, out of 10 people with diabetes, most of them said that they came to Puskesmas only to get a free check-up regarding their blood glucose levels. In fact, Puskesmas already had got programs for people with Diabetes mellitus once a month in the second week, including education and exercise. However, in reality, not many people with Diabetes attended the programs. Mostly the sufferers reasoned that they had to take care of their grandchildren at home and they were still busy at works. The fact shows that lack of physical activity can be an obstacle in controlling blood glucose levels.

High blood glucose levels are the cause of diabetes mellitus, which is associated with metabolic problems. Insulin, a hormone secreted by the pancreas, regulates blood glucose levels by allowing cells to absorb sugar from the blood. Diabetes, on the other hand, is characterized by insulin insufficiency, which is produced by a lack of insulin secretion as well as insulin resistance to its receptors (6). In type 2 diabetes mellitus, the pancreas can still produce insulin, but the quality of insulin is poor and cannot put glucose into cells. As a result, glucose in the blood increases (7). Diabetes Mellitus requires continuous medical care and counseling for self-management to prevent acute and chronic complications (8).

Diabetes Control and Complication Trial (DCCT) shows that good diabetes mellitus control can reduce 20-30% of chronic complications of Diabetes Mellitus. When underestimated, the complication of Diabetes Mellitus can cause damage to organ functions, multi-organ failure, especially the eyes, kidneys, heart, nerves, and other blood vessels (8). Thus, the crucial thing needed is DM control through four pillars of diabetes mellitus control guidelines, which consist of education, dietary management, exercise, and medication adherence (2). DM sufferers who are overweight or not actively doing activities are most likely not able to control their blood glucose levels. Walking for 30 minutes can also improve glucose control, which can help muscles absorb blood glucose and prevent blockages in blood flow. These effects can last for hours or days, but they are not permanent. Thus, to control blood glucose, walking exercise should be done regularly (9). Diabetes Mellitus must be treated because the disease can cause various that can be fatal, such as heart disease, impaired

kidney functiocomplicationsn, blindness, leg rot which sometimes requires amputation, and impotence (10) (11).

As a counselor, i.e., a health nurse in the community, a nurse is often asked by individuals and families to solve the health problems at hand. The nurses' roles as educators and counselors play an important role in building self-confidence by optimizing or increasing the confidence of people with diabetes to be able to control the disease. Based on the description above, the authors were interested in examining the effectiveness of walking to reduce blood glucose levels in people with Diabetes mellitus at Puskesmas Kebonsari, Surabaya.

MATERIALS AND METHODS

This research used the time series design method and was conducted in May 2019. The research participants were 15 randomly selected patients with Type 2 Diabetes Mellitus at Puskesmas (public health center) Kebonsari Surabaya. The inclusion criteria were patients aged 45-65 years, blood glucose levels >100 mg/dl, taking OAD (Anti-Diabetes Medicine) drugs. Patients with diabetic foot ulcers were excluded from this study.

The research instrument was a structured questionnaire (sociodemographic data) consisting of 5 question items that involved gender, age, latest educational background, occupation, and duration of suffering from Diabetes mellitus. The instrument for walking interventions used a standard walking operational procedure: a stopwatch to measure the duration of the walk. The instrument for examining blood glucose level was a set of blood glucose checks (i.e., easytouch glucometer, blood glucose test strips, lancet, alcohol swab, and observation sheets).

Interventions for walking were carried out three times in 2 weeks; twice in week 1 (day 1 and day 3) and once in week 2 (day 8), each for 30 minutes. The walking location was the field around Puskesmas, researchers prepared and cleaned the location to minimize the occurrence of injuries. Patients are advised to use closed footwear that is soft and comfortable for walking. Walking therapy is done together in the morning at 7 am.

Data collection was carried out after the researchers obtained permission from the city's National and Political Unity Agency as well as the research location. The researcher gave a questionnaire to measure the sociodemographic status after the patient gave informed consent as a research respondent. The intervention of walking for 30 minutes was carried out on day 1, 3 and 8. Measurement of blood glucose levels was carried out before and after walking. Furthermore, the researchers analyzed the data on the results of the completed questionnaire and data on blood glucose levels. Blood glucose levels were tested by the Shapiro-Wilk normality test >0.05. All data were normally distributed. Data were

analyzed using paired t-test to determine differences in blood glucose levels before and after walking. The initial blood glucose levels (pre-test on day 1), blood glucose levels after walking (day 3 and day 8) were analyzed using repeated ANOVA tests to identify differences in the three measurements of blood glucose levels (CI 95%, $\alpha = 0.05$).

This current research had received ethical approval from the Health Research Ethics Committee of Sekolah Tinggi Ilmu Kesehatan Hang Tuah Surabaya, Indonesia, with a Certificate of Ethical Eligibility Number: PE/31.a/V/2019/KEPK/SHT, dated 8th May, 2019.

RESULTS

The research was conducted on 15 respondents with Diabetes mellitus in the working area of *Puskesmas Kebonsari Surabaya*. Most of the respondents were female (60%), aged 51-60 years (40%), with last educational background of senior high school (SMA) (73.3%), and on average, self-employed working (46.7%). Concerning the history of Diabetes Mellitus, the respondents, on average, had suffered from the disease for 3-5 years (46.7%). The characteristics of respondents can be seen in table I.

Table I : The characteristics of Respondents (n=15)

Characteristics of Respondents		Frequency (%)
Sex	Male	6 (40.0%)
	Female	9 (60.0%)
Age	45-50 years	4 (26.7%)
	51-60 years	6 (40.0%)
	61-65 years	5 (33.3%)
Last educational background	Elementary School	1 (6.7%)
	Junior High School	3 (20.0%)
	Senior High School	11 (73.3%)
Occupation	Homemakers	4 (26.7%)
	Self-employed	7 (46.7%)
	Not working	4 (26.7%)
Duration of suffering from DM	0-2 years	3 (20.0%)
	3-5 years	7 (46.7%)
	6-8 years	5 (33.3%)

Source : primary data

Table II shows differences in blood glucose levels before and after walking on the 1st, 3rd, and 8th days ($p < 0.001$). This study found significant differences in all blood glucose levels measurements of Diabetic Mellitus both at the beginning (before walking therapy), and after walking therapy on the 3rd and 8th day ($p < 0.025$). This means that walking is effective in lowering blood glucose levels of people with Diabetes mellitus (Table III).

Table II : Difference of Glucose levels of the respondents before and after walking therapy (n=15)

Glucose Level (GL)	N	Mean \pm SD	p
GL pre day-1 (before walking exercise)	15	241.60 \pm 36.56	<0.001
GL post day-1	15	215.20 \pm 30.89	
GL pre day-3	15	247.80 \pm 29.71	<0.001
GL post day-3	15	233.47 \pm 29.82	
GL pre day-8	15	242.27 \pm 21.05	<0.001
GL post day-8	15	223.0 \pm 20.06	

paired t-test. Glucose Level (GL)

Table III : Difference of glucose levels of the respondents before and after walking therapy (day-3 and day-8) (n=15)

Glucose levels	Mean \pm SD	p
Initial (before walking therapy)	241.60 \pm 36.56	0.016
Day-3	233.47 \pm 29.82	
Day-8	223.0 \pm 20.06	

Repeated ANOVA test. The test of *post-hoc paired-wise comparison* resulted in the comparison of all groups < 0.025 .

DISCUSSION

Diabetes Mellitus (DM) is a group of metabolic diseases with the characteristic of hyperglycemia that occurs due to abnormalities of insulin secretion, insulin work, or both (1). (2) recommended the need for good DM control to prevent chronic complications. Diabetes is well controlled when blood glucose levels reach the expected level (12). The average blood glucose level of the participants in this study was 241.60 mg/dl, with the lowest being 200 mg/dl and the highest being 302 mg/dl. This data indicate that diabetes control in respondents was very poor.

Factors that affect blood glucose levels include disease and stress, obesity, food intake, physical activity (exercise), treatment both with Oral Hypoglycemic Medications and with insulin, and duration of suffering from DM (7,13). (14) said that 150 patients in general had poor knowledge, attitude and practice towards type 2 diabetes care. The results showed that the majority of sufferers (73.3%) were more than 50 years old (56 \pm 6.85) and most (66.6%) had been suffering from Diabetes > 3 years. This means that as we get older, the body's intolerance to glucose increases, making sugar levels difficult to control.

Various distress conditions will trigger an increase in blood glucose levels, such as the threat of more severe disease (complications of diabetes). Perceptions of high blood glucose, disciplined and regular treatment that must be followed, or ignorance of treatment are various kinds of stressors for people with diabetes. If they are not managed properly, they can trigger an increase in blood glucose. Previous research on 34 DM sufferers showed that most respondents experienced diabetes distress (73.5%), and the dominant factor affecting it was the

threat of more severe disease (increased blood glucose levels with various complications) (16). So it can be assumed that stress and ignorance of diabetes treatment will cause a lack of control of blood glucose levels and can lead to distress conditions that have an impact on rising blood glucose levels.

The first step in managing Diabetes Mellitus that must be done is non-pharmacological management, in the form of meal planning and physical activities (1). This current study found that there were differences in blood glucose levels before and after walking on days 1, 3, and 8 ($p < 0.001$). In this study, the intervention was carried out 3 times, namely twice in week 1 and once in week 2. Before the intervention of walking, the respondent's blood glucose level was measured and an evaluation was carried out after walking.

Periodic evaluation is needed to see the progress of training and to find out the benefits of physical training that has been done. Good and satisfying results will increase the motivation of people with diabetes to continue doing physical exercise (17). The results of this current research showed that, on the 3rd day, the mean blood glucose level after the intervention on foot was 233.47 mg/dl, which means that there was a decrease. Likewise, the mean blood glucose level on day 8 (or week 2) was 223.0 mg/dl, which also indicates a decrease. Measurement of blood glucose levels before and after walking therapy is needed to monitor and prevent the potential risk of hypoglycemia in diabetes pandemic and to evaluate the achievement of walking therapy.

The results also showed a significant difference in all measurements of blood glucose levels in Diabetes Mellitus patients both at baseline (before walking therapy) and after walking therapy on day 3 and day 8 ($p < 0.025$). These data show that walking is effective at lowering blood glucose levels. The efforts of DM sufferers to carry out self-care (Diabetes self-management), both preventing and tackling, will be related to DM control (18).

Taking medication, physical activity, exercise, and eating habits are related to blood glucose levels (19). During exercise, insulin resistance is reduced; conversely, insulin sensitivity increases, thereby reducing insulin requirements in people with type 2 diabetes. This response only occurs every time a person exercises; it is not a permanent or long-lasting effect. Therefore, exercise must be carried out continuously and regularly (17). This study assumes that, in addition to anti-diabetes drugs, blood glucose levels are influenced by the patient's ability to perform self-care, patient compliance in physical activity and sports, and dietary compliance.

Overweight is the most significant risk factor (20). Walking can reduce the blood glucose level. To lower

blood glucose levels, a person can walk for 30-50 minutes per day. Doing the physical activity in everyday life is one way to control the blood glucose levels of DM patients (9). In line with the research of (21), 30 minutes of leisurely walking exercise can reduce blood glucose levels. Physical activity needs to be done not only for people with diabetes but also for individuals with a high risk of type 2 diabetes. They need to start a healthier life by maintaining their physical activity, especially in sports (22). This study argues that walking not only moves the lower extremity muscles but also moves the upper limb muscles. The more active body muscles, the greater the benefits of exercise that a person will get.

This study adds to the evidence that a walking program can lower blood glucose levels. As explained in the previous studies, a walking exercise program can reduce fasting blood glucose levels (23–25), random blood glucose levels (21,26), can increase daily physical activity, physical strength, and energy consumption (behavioral aspects), and reduce HbA1c and triglycerides (biochemical aspects) in people with type 2 diabetes (25). Furthermore, the pedometer-based walking program can lose weight and improve metabolic health such as fasting glucose, cholesterol, and blood pressure (27).

In this study, the walking intervention was carried out 3 times, namely twice a week with a duration of 30 minutes for each activity. It was done so considering that walking is an activity and physical exercise that is cheap, without any additional tools, and easy to do. In addition, most of the respondents in this study were aged 50 years and over. People with diabetes are recommended to adopt a healthy lifestyle so that blood glucose levels can always be under control, for example by adjusting their diet and increasing physical activity (e.g., exercising or another physical exercise). Physical exercise can increase blood flow. More capillaries open, resulting in the number of insulin receptors so that the receptors will be more active and have an impact on reducing blood glucose levels in people with diabetes (1).

The limitation of this study is that several confounding variables cannot be controlled by the researchers which may also affect the results of the study, including daily physical activity, diet, and stress in people with diabetes mellitus.

CONCLUSION

Walking is one of the physical exercises that DM people can do to control their blood glucose levels. Walking that is carried out 2 to 3 times a week with a duration of 30 minutes is effective in reducing blood glucose levels in type 2 Diabetes Mellitus. Health education on self-care management, especially physical exercise on walking, will support the control of blood glucose

levels. However, further research is needed regarding pedometer-based walking physical exercise on the blood glucose levels of DM patients by controlling confounding variables.

ACKNOWLEDGMENTS

We would like to express our gratitude to all research respondents with type 2 Diabetes mellitus for their participation in this study. We would also like to express our gratitude and high appreciation to the entire academic community of Sekolah Tinggi Ilmu Kesehatan Hang Tuah Surabaya for their continuous support.

REFERENCES

- Soegondo S, Soewondo P, Subekti I, editors. *Penatalaksanaan Diabetes Mellitus Terpadu*. Jakarta: Fakultas Kedokteran Universitas Indonesia; 2015.
- PERKENI. *Konsensus Pengendalian dan Pencegahan Diabetes Melitus Tipe 2 di Indonesia 2011* [Internet]. Perkeni. PERKENI; 2011. 78 p. Available from: https://www.labcito.co.id/wp-content/uploads/2015/03/Revisi_Final_KONSENSUS_DM_Tipe_2_Indonesia_2011.pdf
- World Health Organization. *Diabetes* [Internet]. 2021 [cited 2021 May 25]. p. 1. Available from: <https://www.who.int/news-room/fact-sheets/detail/diabetes>
- Kementerian Kesehatan Republik Indonesia. *Cegah, Cegah, Dan Cegah: Suara Dunia Perangi Diabetes* [Internet]. 2018 [cited 2021 May 25]. p. 1. Available from: <https://www.kemkes.go.id/article/view/18121200001/prevent-prevent-and-prevent-the-voice-of-the-world-fight-diabetes.html>
- Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan RI. *Laporan Nasional RISKESDAS 2018* [Internet]. 2019 [cited 2021 May 25]. Available from: https://labmandat.litbang.kemkes.go.id/images/download/laporan/RKD/2018/Laporan_Nasional_RKD2018_FINAL.pdf
- Handaya Y. *Tepat dan Jitu Atasi Ulkus Kaki Diabetes*. 1st ed. Jakarta: Rapha Publishing; 2016.
- Tandra H. *Life Healthy With Diabetes*. Yogyakarta: Rapha Publishing; 2013.
- Putri NHK, Isfandiari MA. Hubungan Empat Pilar Pengendalian DM Tipe 2 dengan Rerata Kadar Gula Darah. *J Berk Epidemiol* [Internet]. 2013 [cited 2018 Nov 25];1(2):234–43. Available from: <http://journal.unair.ac.id/filerPDF/jbed89640f867full.pdf>
- Muhammad A. *Jangan Abaikan Jalan Kaki*. Jogjakarta: DIVA Press; 2011.
- Mahendra et al. *Care Yourself, Diabetes Melitus*. Jakarta: Penebar Plus; 2008.
- Rahmawati FD. Hubungan Tingkat Self Care dengan Kadar HbA1c Pada Pasien Diabetes Melitus tipe 2 di Poli Interna RSUD Sidoarjo. *STIKES Hang Tuah Surabaya*; 2016.
- Pranoto A. *Terapi Insulin Pada Penderita Diabetes Melitus Rawat Jalan Dan Rawat Inap*. Surabaya: Airlangga University Press; 2012.
- Qurratuaeni. Faktor-faktor yang berhubungan dengan ter kendalinya kadar gula darah pada pasien diabetes melitus di RSUP Fatmawati Jakarta tahun 2009 [Internet]. Jakarta: Uin Syarif Hidayatullah.Fakultas Kedokteran Dan Ilmu Kesehatan; 2010 [cited 2018 Nov 26]. Available from: <http://repository.uinjkt.ac.id/dspace/handle/123456789/597>
- Alammy MA, Abdel A, Mohamed N, Mohamed AA. Knowledge Attitude and Practice Among Patients With Type 2 Diabetes Mellitus: a Cross-Sectional Descriptive Study. *Malaysian J Nurs*. 2017;9(1):31–7.
- Aini LMAN. *Asuhan Keperawatan Pada Sistem Endokrin Dengan Pendekatan NANDA NIC NOC*. Jakarta: Salemba Medika; 2016.
- Yuliasuti C, Yuana. Stress in Patients with Diabetes Mellitus. In: *Surabaya International Health Conference* [Internet]. 2017 [cited 2018 Nov 25]. p. 461–8. Available from: <http://journal.unusa.ac.id/index.php/sihc/article/view/353/367>
- Ilyas E. *Latihan Jasmani bagi Penyandang Diabetes Mellitus dalam: Penatalaksanaan Diabetes Mellitus Terpadu*. Jakarta: Balai Penerbit FK UI; 2011.
- Ditjen PPL Depkes RI. *Pedoman Pengendalian Diabetes Melitus dan Penyakit Metabolik* [Internet]. 2008 [cited 2018 Nov 26]. Available from: <https://id.scribd.com/doc/99496432/pedoman-diabetes-depkes-2010>
- Anani S, Udiyono A, Ginanjar P. Hubungan antara Perilaku Pengendalian Diabetes dan Kadar Gula Darah Pasien Rawat Jalan Diabetes Melitus (Studi Kasus di RSUD Arjawinangun Kabupaten Cirebon). *J Kesehat Masy* [Internet]. 2012;1(2):466–78. Available from: <http://ejournal-s1.undip.ac.id/index.php/jkm/article/view/1136>
- Nugroho PS, Tianingrum NA, Sunarti S, Rachman A, Fahrurrozi DS, Amiruddin R. Predictor Risk of Diabetes Mellitus in Indonesia, based on National Health Survey. *Malaysian J Med Heal Sci* [Internet]. 2020 [cited 2021 May 25];16(1):2636–9346. Available from: https://medic.upm.edu.my/upload/dokumen/2020011612061519_MJMHS_0230.pdf
- Widiya AA, Jatmiko SW, Widyatmoko S. Pengaruh Olahraga Jalan Santai Terhadap Kadar glukosa Darah Pada Pasien Diabetes Melitus. *Biomedika*. 2015 Feb 2;7(1).
- Utami GT, Woferst R, Lubis SL. An overview of physical activities among family members with risk of type 2 diabetes mellitus in Pekanbaru. *Enferm Clin*. 2019 Mar 1;29:26–9.
- Colberg SR, Hernandez MJ, Shahzad F. Blood glucose responses to type, intensity, duration, and timing of exercise. *Diabetes Care* [Internet]. 2013 Oct 1 [cited 2021 May 24];36(10):e177–e177.

- Available from: www.BigBlueTest.org
24. Paula TP, Viana L V., Neto ATZ, Leitro CB, Gross JL, Azevedo MJ. Effects of the DASH Diet and Walking on Blood Pressure in Patients With Type 2 Diabetes and Uncontrolled Hypertension: A Randomized Controlled Trial. *J Clin Hypertens*. 2015 Nov 1;17(11):895–901.
 25. Sung K, Bae S. Effects of a regular walking exercise program on behavioral and biochemical aspects in elderly people with type II diabetes. *Nurs Heal Sci* [Internet]. 2012 Dec 1 [cited 2021 May 24];14(4):438–45. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1442-2018.2012.00690.x>
 26. Tasman. Pengaruh Latihan Fisik Jalan Kaki Terhadap Penurunan Kadar Gula Darah Pada Pasien Diabetes Mellitus Tipe II Di Kota Padang. *Menara Ilmu* [Internet]. 2017 Oct 16 [cited 2021 May 25];2(77). Available from: <https://jurnal.umsb.ac.id/index.php/menarailmu/article/view/393>
 27. Fauzi NFM, Mokhtar R, Halim HA, Zailani MH, Isa A. A 10-Week Pedometer-Based Walking Program Induced Weight Loss and Improved Metabolic Health in Community-Dwelling Adults. *J Sains Kesihat Malaysia*. 2019 Jan;17(01):21–9.